



Arkansas Statewide Collegiate Substance Use Assessment, 2024



Arkansas Collegiate Substance Use Assessment 2024

Derek R. Slagle, Ph.D.

Adam Williams, Ph.D.

Jessica Amos-Dolls, M.P.A.



**ARKANSAS
DEPARTMENT OF
HUMAN
SERVICES**

**Office of Substance
Abuse and Mental Health**

ABOUT THIS REPORT:

Sponsored by the University of Arkansas at Little Rock MidSOUTH Center for Prevention and Training.

Funded by Arkansas Department of Human Services, Office of Substance Abuse and Mental Health (OSAMH) with continued support from the Substance Abuse and Mental Health Services Administration's Center for Substance Abuse Prevention (SAMHSA/CSAP) via the Substance Abuse Block Grant.



MidSOUTH CENTER FOR
PREVENTION AND TRAINING
UNIVERSITY OF ARKANSAS AT LITTLE ROCK

Research coordination and management of the assessment facilitated through the University of Arkansas at Little Rock Survey Research Center

Use of the information contained in this report is encouraged. We require that use of any kind be acknowledged through appropriate citation; for example, we suggest the following format: *Slagle, D.R., Williams, A., & Amos, J. (2025). 2024 Arkansas Collegiate Substance Use Assessment. Little Rock, AR: UA Little Rock Survey Research Center, University of Arkansas Little Rock.*

Short Reference: UA Little Rock Survey Research Center (2025) 2024 Arkansas Collegiate Substance Use Assessment.



For print, broadcast, and other media (radio or television broadcasts, and social media, or other electronic outlets, internet, weblogs, Twitter, Google+, etc.), we suggest an acknowledgment as follows: *The information is based on a report by the University of Arkansas Little Rock Survey Research Center.*

Table of Contents

KEY FINDINGS:	7
INTRODUCTION	9
OVERVIEW:	9
LITERATURE REVIEW:	9
IMPORTANCE OF ASSESSMENT:	9
RISK FACTORS SPECIFIC FOR SUBSTANCE USE IN COLLEGE CAMPUSES:	10
MENTAL HEALTH ISSUES IN COLLEGE CAMPUSES:	11
ALCOHOL:	12
TOBACCO & VAPING:	16
MARIJUANA:	23
SUBSTANCE USE:	24
PRESCRIPTION DRUGS:	25
METHODOLOGY:	26
SURVEY INSTRUMENT AND DISTRIBUTION:	26
STUDENT SELECTION:	27
RESPONDENT CHARACTERISTICS:	29
RESULTS	33
RISK BEHAVIORS	33
ALCOHOL	37
<i>AGE OF FIRST USE</i>	<i>37</i>
<i>LIFETIME USE</i>	<i>38</i>
<i>RECENT USE</i>	<i>39</i>
TOBACCO	42
<i>AGE OF FIRST USE</i>	<i>42</i>
<i>LIFETIME USE</i>	<i>43</i>
<i>RECENT USE</i>	<i>44</i>
VAPING	46
<i>AGE OF FIRST USE</i>	<i>46</i>
<i>LIFETIME USE</i>	<i>47</i>
<i>RECENT USE</i>	<i>48</i>
MARIJUANA	52
<i>AGE OF FIRST USE</i>	<i>52</i>

<i>LIFETIME USE</i>	53
<i>RECENT USE</i>	54
<i>PRESCRIPTION MEDICATION</i>	55
<i>AGE OF FIRST USE</i>	55
<i>LIFETIME USE</i>	57
<i>RECENT USE</i>	58
<i>ILLEGAL DRUGS</i>	60
<i>AGE OF FIRST USE</i>	60
<i>LIFETIME USE</i>	64
<i>RECENT USE</i>	68
<i>CAMPUS OBSERVATIONS</i>	73
<i>POLICIES AND PROGRAMS</i>	73
<i>CAMPUS ENVIRONMENT AND STUDENT SUBSTANCE USE</i>	74
<i>MENTAL HEALTH INDICATORS</i>	80
<u>REFERENCES:</u>	86

Key Findings:

The 2024 survey data reveal significant insights into substance use patterns among college students. For alcohol consumption, most students reported never experiencing alcohol-related problems in the past 30 days, with over 80% indicating no instances of hangovers, poor performance, police involvement, missed classes, fighting, property damage, DUI incidents, criticism, or nausea/vomiting. The data shows important age-related patterns, with those under 21 more likely to abstain from alcohol compared to their older peers, though a substantial percentage of underage students reported some alcohol consumption.

Regarding tobacco use, approximately 64% of respondents reported never using tobacco products, with significant differences between age groups. Students under 21 showed much higher abstinence rates (78%) compared to those 21 and older (55%). Similarly, e-cigarette and vaping products showed notable usage patterns, with about 67% reporting never using these products. Interestingly, the data revealed vaping as particularly prevalent, with nearly 42% of students perceiving daily vaping among their peers—the highest perceived daily use rate among all substances surveyed.

For marijuana, about 56% reported never using the substance, with age again proving to be an important factor. Students under 21 were more likely to report never using marijuana (67%) compared to those 21 and older (48%). When examining prescription drug misuse, most students (84-90%) reported never using prescription opioids or others' prescription medications, with even higher abstinence rates among younger students.

The survey data also revealed campus perceptions, with over 96% of students aware of campus alcohol policies and nearly 91% believing these policies are enforced. Students widely recognized campus concerns about substance use (90%) though relatively few (17%) reported personal involvement in campus prevention programs. Mental health indicators showed significant prevalence of emotional distress among students, particularly feelings of nervousness, depression, and hopelessness, suggesting important connections between mental health and substance use patterns.

Demographically, the survey sample was predominantly female (74%), white (68%), single (69%), and full-time students (82%), with a notable portion being first-year undergraduates (34%). Most reported good academic standing with 90% maintaining A or B grade averages, providing context for understanding substance use patterns within this population.

General Findings:

- The survey encompasses 2-year, 4-year, and post-baccalaureate institutions, as well as public and private universities & colleges.
- The 16-20-year-old age range is repeatedly the age where respondents identified first use of alcohol, tobacco, and other drugs.
- Across the demographics surveyed, most of the participants in this survey study were White (68%) Female (74%), single (69%), full-time (82%), in-state residents (86.5%), with a notable portion being first-year undergraduate student status (34%).
- Regarding student status, most students were staying off-campus (66.5%).
- For class modality, most students were taking hybrid classes (40%).
- Risk behaviors were consistent, albeit somewhat decreased, from the previous post-COVID assessments.
- Nearly 50% of respondents noted a family member with a substance use history.

Alcohol:

- Across all 4 years of assessments, the highest incident age of first alcohol usage among survey participants was 16-20.
- Lifetime occasions of alcohol use was higher for 21+ (legal drinking age) vs. 18-20; similarly, 26+ have a higher lifetime occasions of alcohol use when compared to 18-25.
- When asked about alcohol consumption in the last 30 days, most of the participants had never been drunk or been high from drinking alcoholic beverages (79%).
- When asked about >5 + drinks in a sitting, many of the survey participants who had never had >5 + drinks in a sitting over the last 2 weeks was 84%.

Tobacco & Vaping:

- While most respondents had never used tobacco or vaped, the highest incidence for age of first tobacco usage and vaping was 16-20 years old.
- Lifetime occasions of tobacco use, and vaping was higher for 21+; similarly, 26+ have a higher lifetime occasions of

tobacco usage use when compared to 18–25-year-old. However, there were greater incidence of 18-25 vaping than 26+.

- Majority of the participants (84%) had never used tobacco or vaped in the past 30 days.
- Most participants who reported vaping nicotine

Marijuana:

- Between the ages of 16-20 was the highest incidence of first-time using marijuana at 26% of respondents.
- Recent Marijuana Use was 15% of respondents.

Substance use and Prescription drugs abuse:

- Majority of the participants had never used prescription opioids (84%) or another person's prescription drugs (90%).
- Lifetime occasions of substance use (10+ times usage) was seen to be highest for Amphetamines (7.2%). More than 95% of the participants had never used a substance such as cocaine, amphetamine, sedatives, hallucinogens, opiates, designer drugs, steroids, meth, and illegal drugs .

Campus perceptions:

- Most of the participants (97%) were aware of the campus policies/programs for alcohol, tobacco and substance use and most (91%) believed that they were enforced as well. However, 83% of respondents were not involved in campus prevention efforts.
- Many participants thought that the social atmosphere on the campus does not promote alcohol (78%) or any other drug use (87%).
- Relative to other campuses in the state, close to half (50%) of the participants thought that their campus use was lower than campuses they were familiar with.

Mental Health Indicators:

- Having recent feelings (<30 days) of:
 - Nervousness (74%)
 - Hopelessness (42%)
 - Restlessness (63%)
 - Worthlessness (35%)
 - Depression (51%)
 - Helplessness (57%)

INTRODUCTION

Overview:

The fourth annual statewide collegiate substance use assessment was conducted by the University of Arkansas at Little Rock Survey Research Center from August to December 2024 using a web-based survey instrument. The assessment was sponsored by the University of Arkansas at Little Rock MidSOUTH Center for Prevention and Training and funded by Arkansas Department of Human Services Office of Substance Abuse and Mental Health with continued support from the Substance Abuse and Mental Health Services Administration's Center for Substance Abuse Prevention (SAMHSA/CSAP) via the Substance Abuse Block Grant. The purpose of the assessment was to collect self-reported information on college students' substance use, the behaviors and consequences related to use, and students' perceptions of substance use on college campuses across the state of Arkansas. The study, now having completed its fourth year, has collected 11,652 responses from 52 different colleges and universities across the state of Arkansas. Respondents span the scope of student and life experience and provide a robust sample for analyzing substance use, mental health outcomes, and campus implications.

LITERATURE REVIEW:

Importance of Assessment:

The Arkansas Epidemiological Outcomes Work group indicated there is a statewide deficiency of collegiate-level data for Arkansas that measures the incidence of substance use within the college population (AFMC, 2019). The collegiate level data is important based on a variety of factors, but this data is particularly relevant given that "this category of the population may present as a high risk for first-time users of illicit substances" (AFMC, 2019, p. 13). The data collected for the Arkansas Collegiate Substance Use Assessment represents the only systematic Arkansas statewide data collection where college students self-reported information related to

substance use, consequences of use, and perceptions of substance use on college campuses. Data collected as part of this assessment will be utilized to create safer campuses and will leverage the Substance Abuse Block Grant, which is set to help plan, implement, and evaluate activities that prevent and treat substance abuse on Arkansas college campuses.

Risk Factors Specific for Substance Use in College Campuses:

College enrollment represents a distinct phase in life that initiates the transition from adolescence to independent adulthood and a period of increased exposure to the pressures of alcohol, tobacco, and other substance use (Welsh, Shentu, & Sarvey, 2019). Over the past decade there has been a nationwide increase of cannabis, stimulant, and other illegal drug use across U.S. college campuses (Welsh, Shentu, & Sarvey, 2019) to the point that substance use disorder has become one of the most pervasive health problems for American college campuses (Schulenberg, Johnston, O'Malley, & Bachman, 2017). Substance use disorder is also a major behavioral and health concern among the college students themselves (Rimsza & Moses, 2005). Substance use is associated with multiple negative outcomes for college students including higher probability of unemployment after graduation, lower academic performance, failure to graduate, and increased risk of committing or experiencing sexual assault (Arria, Caldeira, Bugbee, Vincent, & O'Grady, 2015; Arria, et al., 2013; Horsman, 2014; Rimsza & Moses, 2005; Wolaver, 2002;), as well as associations with significant general medical and psychiatric mortality and morbidity for some students (Skidmore, Kaufman, & Crowell, 2016; White, Hingson, Pan, & Hsiao-ye yi, 2011). There is an added importance for higher education institutions to address substance use among students given the unique nature of planning treatments for colleges (e.g., aspects of confidentiality, unique financial constraints, potential university involvement/oversight) (Welsh, Shentu, & Sarvey, 2019). Additional risk factors specific to college campuses and the collegiate population are included in the narratives for the different substances as presented within the report.

Mental Health Issues in College Campuses:

The escalating concerns surrounding mental health on college campuses in the United States demands attention as students grapple with issues like academic pressure, peer expectations, the transition to college life, and societal changes, resulting in prevalent disorders such as anxiety, depression, and stress. University student mental health and wellness has become an important public health concern (Brown, 2016). Hunt and Eisenberg's study in 2010 emphasized the significance of addressing these issues promptly, revealing a substantial proportion of students experiencing emotional distress. The WHO World Mental Health International College Student project (see Auerbach et al. in 2018) underscored that a noteworthy percentage of first-year college students in multiple countries, including the United States, grapple with common mental disorders and that the need for mental health services surpasses the current available resources, emphasizing the magnitude of the situation and the necessity for comprehensive reinforcement strategies. Research has indicated that at least 35% of assessed collegiate students met diagnostic criteria for at least one mental health condition (Auerbach, et al, 2016). When compared to the general population, university and college students are 12.9% higher for levels of depression and age-matched peers (Eisenberg, Gollust, Golberstein, & Hefner, 2007; Ibrahim, Kelly, Adams, & Glazebrook, 2013; Lim, et al., 2018; Cvetovski, Reavley, & Jorm, 2012). The transition from high school to university and developmental transitions to adulthood, prior to 24 years of age are peak onset periods for mental health problems (Sheldon, et al., 2021; Kessler, et al., 2007). Mental health issues have been associated with poor academic performance and higher attrition rates (Hysenbegasi, Hass, & Rowland, 2005; Megivern, Pellerito, & Mowbray, 2003).

Furthermore, the global impact of the COVID-19 pandemic has exacerbated mental health challenges among students worldwide, including in the USA. Wang et al. (2020) highlighted elevated levels of anxiety and depression among college students during the pandemic, adding a

layer of complexity to mental well-being on campuses. The COVID-19 pandemic, and resulting economic and social impacts, have led to global widespread adverse psychological issues, depression, and anxiety (Qiu, et al., 2020) which have been tied to increased substance use and increased addictive/behavioral addictions (Alexander & Ward, 2018). The COVID-19 pandemic and various public health control measures may have increased risks associated with addictive behaviors and substance abuse (Sun, et al., 2020; National Institution on Drug Abuse of USA, 2021; Sun, Bao, & Kosten, 2020). Previous studies on the impacts of disasters have shown higher rates of alcohol use, smoking, and increased risk behaviors (DiMaggio, Galea, & Li, 2009; Lee, Kang, Bell, & Marmot, 2014). Within the college-aged population, a recent study (Firkey, Sheinfil, & Woolf-King, 2021) indicated that domestic college students reported a 26.9% increase in alcohol consumption and a 15.1% increase in cannabis use in response to the COVID-19 pandemic. Concomitantly, recent provisional data (2021) from the Center for Disease Control and Prevention's National Center for Health Statistics demonstrated that overdose deaths in the United States hit an all-time high for the 12-month period ending in 2021 – which was a 28.5% increase from the previous year. This dramatic increase in overdose deaths was seen in opioids, synthetic opioids, methamphetamines, cocaine, and natural and semi-synthetic opioid (prescription pain medication).

Alcohol:

Alcohol and other substance use broadly, have become part of the normative tradition, and integrated into the 'fabric of the college experience' (Welsh, Shentu, & Sarvey, 2019). In fact, full-time college students tend to consume more alcohol than others in their respective age groups (Substance Abuse and Mental Health Services Administration, 2021) and alcohol contributes to an estimated 1,519 deaths annually for college students (Hingson, Zha, & Weitzman, 2009). Similarly, over the past couple of decades, it has been established that college students are at a

higher risk for binge-drinking than non-college students (Johnston, O'Malley, & Bachman, 2011). The typical student embarks on their educational journey two to three years before reaching the legal drinking age, which means that alcohol becomes a legal and consumable substance for them as they are likely entering their final year of school. The transition to the legal drinking age on college campuses creates a policy-spectrum of rules that can create confusion for those students that are subject to the enforcement policies (Cremeens et al., 2011).

The overlap between this annual statewide collegiate assessment and the annual statewide high school assessment is worth observing because there is an observed relationship where students bring established drinking habits from high school to the college level (Schulenberg et al., 2017). Additionally, as college students near the legal drinking age of 21 there is an increased risk for an alcohol use disorder (SAMHSA, 2021). The prevalence of such alcohol use disorders among college students rose from 104,000 students for 18-year-olds to more than double that number, 231,000 by the age of 21 (U.S. Department of Health and Human Services, Substance Abuse and Mental Health Services Administration, 2020). Historically, drinking by college males has exceeded female alcohol consumption but this trend has reversed over time and the most recent 2019 data indicated that females are consuming more alcohol than their male counterparts – however males are still binge drinking and heavy drinking at higher rates (SAMHSA, 2021).

The Substance Abuse and Mental Health Services Administration (2021) reported that in 2019, 53% of full-time college students drank alcohol in the past month (33% of which reported binge drinking) and 8% reported heavy drinking in the past month. Impairment and elevated blood alcohol concentrations associated with binge drinking place those individuals, and individuals around them, at a significantly elevated risk for negative consequences (e.g., injury-related deaths, traffic accidents, sexual assault, violent crimes, and poor academic performance) (U.S. Department

of Health and Human Services, Substance Abuse and Mental Health Services Administration, 2018), and increased illicit drug use (O'Grady, Arria, Fitzelle, & Wish, 2008). A quarter of community-college students engaged in binge-drinking (heavy episodic drinking – typically defined as more than 5 drinks in 1 sitting) and community college student binge drinkers had higher rates of drinking-related problems than non-binge students or abstainers (Sheffield, Darkes, Del Boca, & Goldman, 2010). About one-in-four college students reported having negative consequences associated with drinking (including falling behind in coursework, missing classes, doing poorly on exams/reports, and overall receive lower grades) (White & Hinson, 2013). There are associations between alcohol use and deaths, unintentional injuries, physical fights, and other substance abuse (National Institute of Alcohol Abuse and Alcoholism, 2002; Wechsler, Lee, Kuo, & Lee, 2000; Clements, 1999; Gledhill-Hoyt, Lee, Strote, & Wechsler, 2000) Binge drinkers are more likely to miss class and get hurt related to their alcohol use and can negatively impact other non-drinking students across campus (Wechsler, Lee, Kuo, & Lee, 2000).

The Arkansas Collegiate Substance Use Assessment included a question on binge drinking. The National Institute on Alcohol Abuse and Alcoholism (NIAAA) defined binge drinking as, “a pattern of drinking alcohol that brings blood alcohol concentration (BAC) to 0.08 percent—or 0.08 grams of alcohol per deciliter—or higher” (NIAAA, 2021, What is Binge Drinking, para. 1). The NIAAA further explains that “[f]or a typical adult, this pattern of alcohol misuse corresponds to consuming 4 or more drinks (female), or 5 or more drinks (male) in about 2 hours” (NIAAA, 2021, What is Binge Drinking, para. 1). Alcohol use on college campuses remains a heavily researched topic due to its direct correlation with the long-term effects on underage individuals (Horsman, 2014). In a recent study, the risk of negative consequences was found to be positively correlated with the consumption of alcohol and cannabis on college campuses (Gunn et al., 2022).

Colleges and universities have implemented environmental approaches to reduce alcohol consumption within college students primarily through enforcement and policy development mechanisms (DeJong & Langford, 2002; Neighbors, et al., 2007). Universities and colleges are mandated by the 1989 Drug Free Schools and Communities Act Amendments to put programs into practice for the prevention of illicit drugs and student alcohol abuse (Neighbors, et al., 2007; DeJong & Langford, 2002; Lipperman-Kreda, Grube, & Paschall, 2010). An analysis of alcohol college policies on campus noted that 61% of campuses proactively enforced alcohol policies; larger universities reported greater enforcement levels; and approximately half noted a need for greater coordination with local law enforcement for on-campus enforcement (Toomey, Miazga, Lenk, Erickson, & Winters, 2011). Institutional size and composition have an impact on the types of policies implemented (Lenk, Erickson, Nelson, Winter, & Toomey, 2012) State and local policies have been indicated to further minimize access, consumption, and alcohol-related problems (Mitchell, Toomey, & Erickson, 2005).

One study of college student awareness of campus policies noted that while 89% of students were aware of campus policies, only 44% of students accepted those regulations and rules and 79% of students drank despite being in violation of campus policies (Marshall, Toberts, Donnelly, & Rutledge, 2011). The finding that college control policies have been limited in their outreach is a well-documented finding (Toomey & Wagenaar, 2002). Additionally, implementing alcohol control policies that regulate the distribution and availability of alcohol can be an effective control mechanism to reduce alcohol use and alcohol related problems (Babor, et al., 2003). Student knowledge of alcohol policies, access to policies (hard-copy or digital), and a student ability to comprehend the policies as written are essential for effective on-campus policies and programming efforts (Marshall, Toberts, Donnelly, & Rutledge, 2011). Research has noted that

alcohol policies on college campuses are often incomplete or difficult to access (Faden & Baskin, 2002) and that the enforcement domain is often of importance and the primary focus (Ringwalt, Paschall, & Gitelamn, 2011) – with states with higher alcohol policy enforcement having fewer alcohol-related arrests, alcohol-related-disciplinary actions, and rapes (Kasimanickam, et al., 2024). Common challenges for college administrators include mixed messages received by students; inconsistency in enforcement, and students’ general attitudes towards alcohol use (Cremeens, et al., 2011). The most common recommendations to address college alcohol policies were ensuring a comprehensive approach with the entire community and enhanced alcohol education (Cremeens, et al., 2011).

Tobacco & Vaping:

Evidence underscores the significant impact of tobacco-related health disparities (TRHDs) on population health in the United States (Berg et al., 2021). Smoking cigarettes, a key contributor to TRHDs, results in nearly 500,000 deaths annually in the United States, with over 40,000 deaths attributed to secondhand smoking. It is noteworthy that almost all adult cigarette smokers initiate smoking before the age of 26 (Wang et al., 2018) and in the general population 99% of adult cigarette smokers began smoking before the age of 26, with many becoming regular, daily smokers during young adulthood. College students experience tobacco-related health disparities like other adult subpopulations (e.g., racial/ ethnic minorities; lower socioeconomic backgrounds/ urban/rural; sexual and gender minorities). Interestingly, tobacco use incidence is higher at community colleges and tribal universities/ colleges -compared to traditional 4-year universities/ colleges (Berg, et al., 2021; McIntosh, Wall, & Johnson, 2016; Lenk, et al., 2012; Choi, Nazir, & Panheco, 2016). Thus, college student populations are a critical target population when attempting to mitigate tobacco-related health disparities (Berg, et al., 2021).

African American and Black populations are more likely to start using tobacco during adulthood rather than in their youth (Roberts, Colby, Lu, & Ferketick, 2016) and those smoking adults have greater morbidity and mortality from smoking-related diseases even though they have lower smoking intensities (Leventhal, Dai, & Higgins, 2022; Cunningham, et al., 2017). African American adults reported higher motivations to quit tobacco yet experienced greater difficulty than other groups due to lack of access to culturally competent care and structural inequities (Carrol & Cole, 2022; Kulak, Cornelius, Fong, & Giovino, 2016). Derefinko, et al. (2018) found that there were no rural/ urban differences broadly in tobacco usage while transitioning to college although rural racial/ ethnic minorities were more likely to use tobacco during these transitions. Historically, tobacco cessation programs have had difficulty recruiting and enrolling minority populations (Coday, Richey, & Thomas, 2016). Black and African American adults demonstrated interest in higher Quitline services and telephone coaching (Marshall, et al., 2017; Webb Hooper, Carpenter, & Salmon, Web-based tobacco cessation interventions and digital inequality across US racial/ ethnic groups, 2019; Zhu, et al., 2011) but have less success than White adults (Rabius, Wiatrek, & McAlister, 2012; Sood, et al., 2009; Varghese, et al., 2014). Therefore, there is a necessity to develop culturally specific tobacco programming and interventional cessation among Black and African Americans to address social norms, racial discrimination/ stressors, menthol smoking, cultural values and beliefs, and history (Hooper, Carpenter, Salmon, & Resnicow, 2023). Culturally focused interventions and programming are preferable to standard, general audience, cessation interventions for Black adults (Webb, 2009; Webb Hooper, Antoni, Okuyemi, Dietz, & Rescinow, 2017).

The prevalence of cigarette smoking and tobacco use hit a historic low in 2020, marking a 4% usage among college students in the past 30 days (Schulenberg, et al., 2020). Despite the

overall decline in tobacco use, the void left has been filled by the rising trend of vaping through electronic cigarettes and vaporizer devices across the United States (Cullen et al., 2019; Miech, Miech, Johnston, O'Malley, Bachman, & Patrick, 2019; Patrick, O'Malley, & Johnston, 2017). The U.S. Centers for Disease Control and Prevention and the Food and Drug Administration demonstrated in the National Youth Tobacco Survey that e-cigarettes/ vaping has been the most common tobacco usage among youth since 2014. Nationwide, approximately 17-40% of all college students have tried or are actively using e-cigarettes (Jones, Asare, & Lanning, 2021). Overall, 60% of college students have been offered an e-cigarette (Copeland, Peltier, & Waldo, 2017; U.S. Department of Health and Human Services, 2016), and, of active users, 40% had not actively used combustible cigarettes (Schoenborn & Gindi, 2016) – despite the fact that e-cigarette use is positively associated with traditional combustible cigarettes in the future (Coleman, et al., 2015). In aggregate, these findings demonstrate that vaping usage/ e-cigarettes are generating new populations of individuals susceptible to developing a nicotine addiction (Jones, Asare, & Lanning, 2021).

The surge in vaping has been linked to various factors such as harm reduction, the absence of regulation, the appeal of public consumption (Trumbo, 2017; Tavollaci, et al., 2016; Franks, Hawes, McCain, & Payakachat, 2017), and directed marketing claims made at college students for vaping (Luo, Zheng, Zeng, & Leischow, 2014). Among targeted advertisements for college e-cigarette use, 95% of the websites had health-related claims, 64% had smoking cessation-related claims, and 22% featured health professionals. These targeted advertisements and broader marketing have led to the perception, among nonusers and users, that vaping is a healthier alternative to cigarettes and reporting that vaping contains less chemicals, fewer nicotine, and less smoke than normal tobacco cigarettes (Case, Crook, Lazard, & Mackert, 2016). College students

routinely cite the belief that vaping is a healthy alternative to tobacco/ combustible cigarette usage as part of the motivation to use (Hefner, Solazzo, Mullaney, Coker, & Sofuoglu, 2019; Lanza & Teeter, 2018).

Despite being perceived as a potentially less harmful method of nicotine consumption, the surge in vaping among adolescents in the United States has become a major cause for concern. Research has indicated that vaping was associated with smoking, poorer mental health, and alcohol consumption among college students (Wilson, Bullen, Duffey, & Bopp, 2022). Associations have been observed with regular tobacco use, concurrent use of tobacco and cannabis, alcohol consumption, and even suicidal ideations (Lee & Lee, 2019; Auf, et al., 2019; Jones, Hill, Pardini, & Meier, 2016; Oh, et al., 2019). Studies suggest further linkages between vaping and beliefs, social cues, cognitive-risk perceptions, and even residence in urban versus rural areas (Trumbo, 2017; Dai & Hao, 2017); gender, certain risk behaviors, and geographic locations of high schools were predictors of vaping in college (Omoike & Johnson, 2020). The trend extends across educational levels, impacting middle school, high school, and college students, with estimates indicating a vaping rate as high as 25% among college students over the past 30 days (Trumbo, 2017). Although there was a leveling off in vaping for 2020, concerning figures persist, such as a 25% annual prevalence of vaping marijuana among college students and a rise in daily use from 5% in 2017 to 12% in 2020 (Schulenberg et al., 2020; U.S. Department of Health & Human Services, National Institutes of Health, 2021).

Research has demonstrated there are variable use patterns across racial/ ethnic groups for youth and college students. One study noted that the most important reason youth vape was for relaxation and stress/ anxiety coping – with racial/ ethnic minorities reporting this vape motivation (Donaldson, et al., 2022). From 2014-2019, across racial and ethnic groups for youth, non-

Hispanic Blacks were the only racial group to increase for dual use and occasional use – while less likely to report frequent use and flavored cigarette use (Dai, Ramos, Faseru, Hill, & Sussman, 2021). Studies have shown associations between experienced discrimination or racism and tobacco use for both combustible tobacco cigarettes and vaping (Agunwamba, et al., 2017; Fahey, Morris, Robinson, & Pebley, 2021). There is a further need for tailored strategies to address disparities in vaping use across racial and ethnic groups (Dai, Ramos, Faseru, Hill, & Sussman, 2021). Ultimately, the persistence of vaping trends, especially among college students, is influenced by the social atmosphere on campuses (Kava et al., 2021) which is why institutional policies and targeted programmatic support is relevant and instrumental in addressing tobacco-related health disparities.

Efforts to address vaping and tobacco usage necessitate the implementation of effective institutional policies as a crucial strategy (Navon et al., 2019), particularly since the persistence of tobacco and vaping trends, especially among college students, is influenced by the social atmosphere on campuses (Kava et al., 2021). United States colleges and university implementation of comprehensive tobacco control policies doubled from 2012-2017 (Wang, et al., 2018). Similarly, the American Nonsmokers' Rights Foundation (2018) documented that of the 6,000 US universities and colleges, 1,909 were 100% tobacco-free; 2,278 were 100% smoke-free campuses; and 1,885 prohibited vaping on school premises (American Nonsmokers' Rights Foundation, n.d.). Another study noted that of 605 colleges/ universities, 54% did not have tobacco or smoke free policies – while 35% had tobacco-free policies and only 10% had smoke-free policies (Trad, et al., 2018). Bayley et al. (2020) found that of institutions adopting policies, 39% of the sample adopted e-cigarette-free policies; 26% had hookah-free policies, and 20% had tobacco-free policies. A 2019 analysis noted that 27% of college students were protected by state laws and institutional

policies at colleges/ universities (Blake, et al., 2018). As of 2015-2017, only three states had 100% smoke-free and/or tobacco-free policies in more than half of their higher education institutions while four states had no known smoke-free and/or tobacco-free campus protections (Blake, et al., 2018) The implementation in tobacco and vaping control policies varies across the various campus types and settings. (Berg, et al., 2021). There has been a specific focus on community/ technical colleges, tribal colleges/ universities and historically Black colleges/ universities with the creation of the Tobacco-Free Campus Initiative in 2012 (McIntosh, Wall, & Johnson, 2016).

The impact of campus policies has been reviewed in the college student population and noted a positive impact for tobacco control policies on campus, including an impact on student tobacco use behavior (Bennett, Deiner, & Pokhrel, 2017); policy impact on reducing smoking behaviors/ prevalence and pro-smoking behaviors (Lupton & Townsend, 2015); and that multicomponent programs resulted in increased initial cessation rates, fewer relapses, and decreased likelihood across campus (Rodgers, 2012). Other studies have noted that campuses without comprehensive policies were more likely to have students smoke on campus, see others smoking, and see others using tobacco products (Ickes, Wiggins, Rayens, & Hahn, 2020). Literature on tobacco control policies noted that institutional bans on smoking decreased prevalence of combustible cigarette smoking (Allen & Stuart, 2019) while vaping/ e-cigarette usage increased following a smoking ban on college campuses (Leavens, et al., 2020). Concomitantly, on-campus tobacco control policies and implementation can impact off-campus tobacco retail availability (Kates, et al., 2016; Barker, Schleider, Ababesh, Johnson, & Henriksen, 2018).

With regard to tobacco users there is a noted policy gap with regard to policy controls (Russette, Harris, Schuldberg, & Green, 2014; Fallin, et al., 2012; Braverman, Geldhof,

Hoogesteger, & Johnson, 2018) and noncompliance mechanisms across campuses has been documented within the literature (Roditis, Wang, Glantz, & Fallin, 2015). Gaps within these compliance mechanisms underscore the further need for monitoring, implementation, and evaluation of tobacco- and smoke-free policies across U.S. campuses. Having clear university communications and policies denoted on tobacco alternatives is also of importance. One study noted that of 581 universities, 1/5th did not specify whether e-cigarettes/ vaping were prohibited (Jun & Kim, 2020). Other major issues for university-based cessation initiatives include appropriateness and effectiveness. A meta-analysis of 14 studies noted that interventions targeted at young adults were more successful in producing positive cessation rates than controls and that interventions targeting adults were also effective at college-aged students (Suls, et al., 2012). Research has also demonstrated that there is a broad student interest in cessation programs across a wide variety of student demographics (e.g., sex, race/ ethnicity, sexual orientation, other sociodemographics) (Berg, Sutfin, Mendel, & Ahluwalia, 2012). Broadly, on-campus student health services are missing the mark with regard to assessing tobacco use specifically and only 1/3rd of students visiting student health centers were screened for usage (Sutfin, et al., 2012) and less than half of student health clinics screen for tobacco across their visits (Sutfin, et al., 2015). There were multiple noted barriers across campus-based student health centers and resources for campus-based cessation (Sutfin, et al., 2012). Beyond campus health facilities, there are promises for programmatic interventions. Research has noted that personalized motivational interventions and face-to-face skills-based interventions were effective for short-term tobacco cessation (Jeffries, et al., 2016). Other technology-based interventions have been found relevant to college student population (Brown, 2013; Gulliver, et al., 2015; An, et al., 2013; Karekla & Savvides, 2019; Simmons, Heckman, Fink, Small, & Brandon, 2013) including text messaging (Mason, Ola,

Zaharakis, & Zhang, 2015; Camenga, Bernstein, Dziura, Fiellin, & Krishnan-Sarin, 2021; Witkiewitz, et al., 2014), social media (Baskerville, Azagba, Norman, McKeown, & Brown, 2016; Ramo, et al., 2018), and incentive-based cessation intervention programs (Berg, et al., 2014; Popp, et al., 2018; Thomas J. L., et al., 2016; Thomas J. L., et al., 2016). There are considerable research gaps with regard to college student programming and tobacco cessation due to lack of control groups; inconsistent outcome measures; small sample sizes; and multi-component programming that don't adequately address all components in analysis (Berg, et al., 2021). Additionally, there isn't sufficient cessation research on tobacco alternatives; the broader campus context; collegiate racial/ ethnic minorities; or college students with mental health issues (Berg, et al., 2021).

Marijuana:

The use of Marijuana has been steadily increasing nationwide and is the third 'most used' substance among Arkansas college students. Schulenberg et al (2017) noted that daily cannabis use among college students doubled between 2007 and 2014. Similarly, the most recent Monitoring the Future report (Schulenberg, et al., 2020), cited that this trend has continued into 2020 where 44% of college students reported using marijuana, 8% used marijuana on a daily/ near-daily basis, and the number of college students vaping marijuana over the past 30 days rose from 5% to 14%. For this assessment, over 12% of respondents aged 18-25 reported using marijuana within the last 30 days. Studies have noted that increased marijuana/cannabis use is important for the college population since the likelihood of use increases in prevalence with successive years in school (Arria, O'Grady, Caldeira, Vincent, & Wish, 2008) and that heavy marijuana use has a short-term impact on learning and memory which can then impair collegiate academic and health outcomes (Arria, Caldeira, Bugbee, Vincent, & O'Grady, 2015; Arria, Caldeira, Bugbee, Vincent,

& O'Grady, 2016). There is a rise in vaping marijuana among college students, with a 25% annual prevalence and a rise in daily use from 5% in 2017 to 12% in 2020.

Substance Use:

“Substance use” is a term that refers to several substances consumed that can potentially cause dependency and detrimental effects on the body (CDC, 2021). Substances within this paper refer to drugs that may lead to dependency that are addressed within the ACSUA. Transitional periods in individuals' lives, such as the period between high school and college, increase the chance of participating in risky behavior (Fromme et al., 2008). For most, college is when individuals are introduced to the world of substances. The transition and the new exposure increase the likelihood that college students will consume substances they may not have consumed before starting college.

The misuse of stimulant medication within the college population is on the rise (Benson, Flory, Humphreys, & Lee, 2015). A meta-analysis and literature review identified academic pressures as the primary driver, often facilitated by obtaining stimulants from peers with prescriptions. In 2020, Schulenberg et al. reported a significant increase in hallucinogen use among college students, with 9% acknowledging usage, nearly double the 2019 rate of 5%. The prevalence of amphetamine usage without a prescription stood at 6.5%, Adderall at 7.2%, and nonmedical use of Ritalin at 1.4%. The Monitoring the Future survey revealed annual prevalence rates for sedatives (1.7%), tranquilizers (2.6%), narcotics other than heroin (1.3%), Vicodin (1.2%), and OxyContin (1.5%), with cocaine use reaching 3.8%.

The escalation in cocaine use is particularly concerning, with a longitudinal study at a large public university (Kasperski et al., 2011) indicated that, by the fourth year of college, 36% of students had been offered cocaine, 13% had used it, and annual prevalence increased from 4% in

Year 1 to 10% in Year 4. Females exhibited more serious patterns of use and a higher likelihood of cocaine dependence. Hallucinogens, MDMA, LSD, and psychedelics are also on the rise, fueled by social pressures and curiosity. In the broader context, a 2020 study reveals that young adults aged 18 to 25 have the highest substance use disorder rate at 24.4% (Johnson, Aug. 2022). Startlingly, 90% of adults with a substance use disorder-initiated use before adulthood (The JED Foundation, 2022). Overdose mortality rates among adolescents aged fourteen to eighteen surged by 94% in one year, from 2.36 deaths per 100,000 in 2019 to 4.57 per 100,000 in 2020, with a further 20% increase from 2020 to 2021 (Johnson, Oct. 2022).

The issue of substance abuse carries multiple adverse outcomes for students, including diminished academic performance, heightened unemployment risks post-graduation, and increased vulnerability to committing or experiencing sexual assault (Welsh et al., 2019). Notably, students with low grade point averages or facing exceptional performance pressures are at a higher risk of misusing prescription stimulant medications like Adderall. Furthermore, it has been observed that female students are more prone to nonmedical stimulant medication use compared to their non-student counterparts (Welsh et al., 2019).

Prescription Drugs:

Even though college students were particularly vulnerable to opioid misuses during the opioid crisis - young adults reported the highest past-year opioid prevalence use for all age groups (U.S. Department of Health and Human Services, Substance Abuse and Mental Health Services Administration, 2017) - there has been a continual 5-year decline for prescription opioid misuses for college students. In 2020, 1% of college students reported non-medical use of opioids within the past year (Schulenberg, et al, 2020). While declining, it is worth noting that opioid use disorders commonly begin during young adulthood and coincide with typical college years/ ages (Johnston,

O'Malley, Miech, Bachman, & Schulenberg, 2016; Hadland, et al., 2017). Research has noted the opioid misuse in college students is associated with several demographic factors and concurring behaviors - such as off-campus living and low cumulative grade point averages (Harries, Lust, Christenson, Redden, & Grant, 2018).

Methodology:

The Arkansas Collegiate Substance Use Assessment (ACSUA) was conducted by the University of Arkansas at Little Rock Survey Research Center from August to December 2024 using a web-based survey instrument through Qualtrics. The assessment was funded by the Arkansas Department of Human Services, Office of Substance Abuse and Mental Health and coordination and management was facilitated through collaboration between the University of Arkansas at Little Rock Survey Research Center (UALR SRC) and University of Arkansas at Little Rock MidSOUTH. The purpose of the assessment was to collect self-reported information on college students' substance use, the behaviors and consequences related to use, and student perceptions of substance use on college campuses across the state of Arkansas.

Survey Instrument and Distribution:

Survey items were adapted from several sources, including the Core Institute Alcohol and Other Drug Database (Southern Illinois University, Carbondale), the Arkansas Prevention Needs Assessment Survey (Arkansas Department of Human Services Office of Substance Use and Mental Health), Texas College Survey of Substance Use (Texas Health and Human Services), and the National Survey on Drug Use and Health (NSDUH) (Substance Abuse and Mental Health Services Administration Center for Behavioral Health Statistics and Quality). This report represents the third assessment of its kind for Arkansas so last year's results are noted as comparisons to be included in the report. A complete list of higher education institutions was developed by the UALR SRC and included both public and private 2-year, 4-year, and

postbaccalaureate institutions. All institutions were invited to participate in the assessment. The method of email distribution and time frame within the August 15th to November 30th period was determined by the individual institutions. The survey uses a convenience sampling methodology, with all Arkansas colleges invited to participate. The participating schools can vary from year to year, based on individual institutions desire to participate. Due to the number of schools participating in the survey and the absence of a true random sampling technique, while results from previous years are occasionally presented, interpreting comparisons across the years should be done with caution. Additionally, the same restraint is advised when generalizing the findings to *all* total Arkansas college students. Reports from previous years' surveys can be found at:

<https://ualr.edu/publicaffairs/survey-research-center/arkansas-college-substance-use-assessments/>

Student Selection:

The selection of students was determined by participating schools – some schools chose random samples to survey, whereas others distributed the assessment to their entire student population. Participation in the survey was completely voluntary at the institutional and participant level. No incentives were provided as in previous years (i.e., random gift card drawing) which impacted total number of responses from the student population. The survey protocol and instrument were approved through the University of Arkansas at Little Rock Institutional Review Board processes. Responses were only recorded upon completion of the assessment. Partial submissions were excluded from the assessment and there were 26 respondents indicated they did not wish to participate or were not at least 18 years of age, which were excluded from analysis. Security measures were included for this year's assessment to prevent multiple submissions; bot detection; relevantID (propriety Qualtrics settings for analysis of respondents' browser; operating

system; & location) to prevent fraudulent responses; and tools to prevent indexing on search engines.

In total, 2,276 students agreed to participate and completed the assessment. In sum there were 30 academic institutions that recorded student responses. Additionally, there were 55 student respondents which did not answer the question identifying the institution and 33 respondents that indicated 'other'.

Respondent Characteristics:


Respondent Demographic Data, 2024

Variable	<i>n</i>	<i>%</i>
Gender		
Male	453	21.38
Female	1573	74.23
Transgender/ Gender Non-conforming	93	4.39
Racial Identity		
Black or African American	442	20.80
Asian	74	3.48
Native American	17	0.80
Alaska Native	0	0.00
Native Hawaiian or Other Pacific Islander	1	0.05
White	1443	67.91
Other	55	2.59
2+ Races	93	4.38
Hispanic/Latino		
Yes	204	9.57
No	1928	90.43
Marital Status		
Single, divorced	159	7.47
Living with domestic partner	111	5.21
Widowed	8	0.38
Single, never married	1478	69.42
Married, and living with spouse	340	15.97
Married, and living separately from spouse	33	1.55
Employed Status		
Yes, part-time	775	36.45
Yes, full-time	570	26.81
No	781	36.74
Military		
Never served in the military	2076	97.33
Only on active duty for training in the Reserves or National Guard	15	0.70
Now on active duty	2	0.09
On active duty in the past, but not now	40	1.88
Residency		
In-state (Arkansas)	1842	86.52

Out of State	245	11.51
Outside the United States	42	1.97
<hr/>		
Student Status		
Full-time (e.g., 12+credits)	1738	81.63
Part-time (e.g., 1-11 credits)	391	18.37
<hr/>		
Degree Status		
Non-degree seeking	73	3.43
Undergraduate	1196	56.26
Associate	484	22.77
Graduate	303	14.25
Other	70	3.29
<hr/>		
Undergraduate Classification		
Freshman (undergraduate)	404	33.84
Sophomore (undergraduate)	281	23.53
Junior (undergraduate)	263	22.03
Senior (undergraduate)	246	20.60
<hr/>		
Graduate Degree		
Masters	182	60.26
Doctorate	81	26.82
Professional Degree	26	8.61
Other	13	4.30
<hr/>		
Campus Living Situation		
On-campus	711	33.47
Off-campus	1413	66.53
<hr/>		
Modality for Coursework		
Fully online courses	513	24.11
Hybrid (both online and in-person courses)	853	40.08
Fully in-person courses	762	35.81
<hr/>		
Estimated GPA by Grade		
A	1165	55.27
B	739	35.06
C	187	8.87
D	12	0.57
F	5	0.24

Note. Due to rounding errors, percentages may not equal 100%.


Frequency of Respondent Age, 2024



Frequency Table for Age Classifications

Variable	<i>n</i>	%
Age Grouping:		
18-24	1399	65.71
25+	730	34.29
Age: Legal Drinking Age		
18-20	861	40.44
21+	1268	59.56

Family Substance History of Respondents, 2024



Respondent Family Substance Use History, 2024

Family Member Identified with Problem	<i>n</i>
Mother	332
Father	548
Stepmother	48
Stepfather	111
Brothers/ sisters	309
Mother's parents	325
Father's parents	287
Aunts/ uncles	573
Spouse	63
Partner	55
Children	23
No Family Drug History	751

Results

Risk Behaviors

Table 1a. Frequency and Percent of Risk Behaviors in the Last 30 Days, 2024

Risk Behavior	<i>n</i>	%
Hangover		
Never	1661	84.06
Once	180	9.11
Twice	74	3.74
3-5 times	39	1.97
6-9 times	9	0.46
10+ times	13	0.66
Poor Performance in Class		
Never	1702	86.22
Once	145	7.35
Twice	69	3.50
3-5 times	42	2.13
6-9 times	8	0.41
10+ times	8	0.41
Trouble with Police or Campus Authorities		
Never	1947	98.63
Once	19	0.96
Twice	4	0.20
3-5 times	1	0.05
6-9 times	0	0.00
10+ times	3	0.15
Missed Class		
Never	1674	85.02
Once	123	6.25
Twice	92	4.67
3-5 times	53	2.69
6-9 times	16	0.81
10+ times	11	0.56
Been in a Fight		
Never	1850	93.72
Once	66	3.34
Twice	29	1.47
3-5 times	17	0.86
6-9 times	3	0.15

10+ times	9	0.46
Damaged Property		
Never	1958	99.24
Once	11	0.56
Twice	2	0.10
3-5 times	0	0.00
6-9 times	0	0.00
10+ times	2	0.10
Driven while Under the Influence		
Never	1880	95.33
Once	34	1.72
Twice	14	0.71
3-5 times	18	0.91
6-9 times	7	0.35
10+ times	19	0.96
Been Criticized for Alcohol and Drug Use		
Never	1887	95.88
Once	30	1.52
Twice	30	1.52
3-5 times	12	0.61
6-9 times	3	0.15
10+ times	6	0.30
Got Nauseated or Vomited		
Never	1680	85.11
Once	159	8.05
Twice	68	3.44
3-5 times	53	2.68
6-9 times	3	0.15
10+ times	11	0.56

Note. Due to rounding errors, percentages may not equal 100%.

Table 1b. Frequency and Percent of Risk Behaviors in the Last 30 Days, 2024

Risk Behavior	<i>n</i>	%
Been Hurt or Injured		
Never	1836	93.77
Once	60	3.06
Twice	35	1.79
3-5 times	17	0.87

6-9 times	4	0.20
10+ times	6	0.31
<hr/>		
Tried to Stop Using Substances		
Never	1817	92.42
Once	70	3.56
Twice	32	1.63
3-5 times	24	1.22
6-9 times	5	0.25
10+ times	18	0.92
<hr/>		
Experienced Memory Loss		
Never	1760	89.39
Once	89	4.52
Twice	40	2.03
3-5 times	44	2.23
6-9 times	10	0.51
10+ times	26	1.32
<hr/>		
Thoughts that May Have a Substance Problem		
Never	1851	93.91
Once	52	2.64
Twice	21	1.07
3-5 times	22	1.12
6-9 times	4	0.20
10+ times	21	1.07
<hr/>		
Have or Attempted to Take Advantage of Another Sexually		
Never	1957	99.59
Once	3	0.15
Twice	2	0.10
3-5 times	0	0.00
6-9 times	1	0.05
10+ times	2	0.10
<hr/>		
Attempted Suicide		
Never	1941	98.83
Once	13	0.66
Twice	4	0.20
3-5 times	3	0.15
6-9 times	0	0.00
10+ times	3	0.15
<hr/>		
Had Suicidal Thoughts		

Never	1828	92.98
Once	65	3.31
Twice	32	1.63
3-5 times	20	1.02
6-9 times	5	0.25
10+ times	16	0.81
<hr/>		
Been Sexually Assaulted		
Never	1920	97.51
Once	25	1.27
Twice	13	0.66
3-5 times	5	0.25
6-9 times	2	0.10
10+ times	4	0.20

Note. Due to rounding errors, percentages may not equal 100%.

Alcohol

Age of First Use

Figure 1. Age of First Use of Alcohol, 2024




Table 2. Age of First Alcohol Use, 2021-2024

Age of First Use	Year			
	2021	2022	2023	2024
Do not use/ Never	973 (29.98%)	967 (25.25%)	278 (24.34%)	553 (26.79%)
Under 10	142 (4.37%)	112 (2.93%)	44 (3.85%)	61 (2.96%)
11-15	484 (14.91%)	715 (18.67%)	200 (17.51%)	356 (17.25%)
16-20	1296 (39.93%)	1462 (38.18%)	500 (43.78%)	842 (40.79%)
21-25	335 (10.32%)	482 (12.59%)	115 (10.07%)	237 (11.48%)
26+	16 (0.49%)	91 (2.38%)	5 (0.44%)	15 (0.73%)

Note. Due to rounding error, percentages may not sum to 100%.

Table 3. Age of First Use of Alcohol by Current Age Groups, 2024

Age of First Use	Age Groups (18-20, 21+)		Age Groups (18-25, 26+)	
	18-20	21 and Older	18-25	26 and Older
Do not use/ Never	351 (42.24%)	200 (16.30%)	447 (31.43%)	104 (16.35%)
Under 10	17 (2.05%)	43 (3.50%)	35 (2.46%)	25 (3.93%)
11-15	140 (16.85%)	216 (17.60%)	222 (15.61%)	134 (21.07%)
16-20	321 (38.63%)	519 (42.30%)	599 (42.12%)	241 (37.89%)
21-25	N/A	235 (19.15%)	118 (8.30%)	118 (18.55%)
26+	N/A	14 (1.14%)	N/A	14 (2.20%)

Note. Due to rounding error, percentages may not sum to 100%.

Lifetime Use

Figure 2. Lifetime Occasions of Alcohol Use, 2024




Table 4. Lifetime Occasions of Use of Alcohol by Current Age Groups, 2024

Lifetime Use	Age Groups (18-20, 21+)		Age Groups (18-25, 26+)	
	18-20	21 and Older	18-25	26 and Older
0 times	326 (39.56%)	167 (13.88%)	412 (29.45%)	81 (12.90%)
1-2 times	103 (12.50%)	65 (5.40%)	137 (9.79%)	31 (4.94%)
3-5 times	95 (11.53%)	84 (6.98%)	148 (10.58%)	31 (4.94%)
6-9 times	86 (10.44%)	76 (6.32%)	133 (9.51%)	29 (4.62%)
10+ times	214 (25.97%)	811 (67.41%)	569 (40.67%)	456 (72.61%)

Note. Due to rounding error, percentages may not sum to 100%.

Recent Use

Table 5. Alcohol Use and Drunkenness in the Last 30 Days, 2021-2024

	Year			
	2021	2022	2023	2024
Alcohol Use				
0 times	1762 (55.48%)	1769 (46.91%)	559 (51.81%)	1139 (56.61%)
1-2 times	712 (22.42%)	858 (22.75%)	270 (25.02%)	475 (23.61%)
3-5 times	356 (11.21%)	552 (14.64%)	126 (11.68%)	239 (11.88%)
6-9 times	142 (4.47%)	291 (7.72%)	68 (6.30%)	0 (0.00%)
10+ times	204 (6.42%)	301 (7.98%)	56 (5.19%)	159 (7.90%)
Drunkenness				
0 times	2528 (79.72%)	2593 (68.85%)	825 (76.60%)	1585 (79.09%)
1-2 times	404 (12.74%)	551 (14.63%)	158 (14.67%)	282 (14.07%)
3-5 times	132 (4.16%)	302 (8.02%)	44 (4.09%)	80 (3.99%)
6-9 times	36 (1.14%)	169 (4.49%)	28 (2.60%)	0 (0.00%)
10+ times	71 (2.24%)	151 (4.01%)	22 (2.04%)	57 (2.84%)

Note. Due to rounding error, percentages may not sum to 100%.

Figure 3. Beverages Consumed in the Last 30 Days, 2024




Figure 3. Drunk in the Last 30 Days, 2024




Table 6. Alcohol Use in the Last 30 Days by Current Age Groups, 2024

Recent Use	Age Groups (18-20, 21+)		Age Groups (18-25, 26+)	
	18-20	21 and Older	18-25	26 or Older
0 times	563 (69.33%)	572 (47.91%)	793 (57.42%)	342 (54.72%)
1-2 times	165 (20.32%)	310 (25.96%)	330 (23.90%)	145 (23.20%)
3-5 times	51 (6.28%)	187 (15.66%)	168 (12.17%)	70 (11.20%)
6-9 times	0 (0.00%)	0 (0.00%)	0 (0.00%)	0 (0.00%)
10+ times	33 (4.06%)	125 (10.47%)	90 (6.52%)	68 (10.88%)

Note. Due to rounding error, percentages may not sum to 100%.

Figure 5. Binge Drinking in the Past 2 Weeks, 2024




Table 7. Binge Drinking in the Last 30 Days, 2021-2024

Frequency	Year			
	2021	2022	2023	2024
None	2759 (84.32%)	2621 (67.66%)	965 (84.13%)	1759 (84.61%)
Once	268 (8.19%)	478 (12.34%)	97 (8.46%)	187 (8.99%)
Twice	124 (3.79%)	355 (9.16%)	40 (3.49%)	75 (3.61%)
3 to 5 times	85 (2.60%)	259 (6.69%)	29 (2.53%)	40 (1.92%)
6 to 9 times	21 (0.64%)	112 (2.89%)	6 (0.52%)	10 (0.48%)
10 or more times	15 (0.46%)	49 (1.26%)	10 (0.87%)	8 (0.38%)

Note. Due to rounding error, percentages may not sum to 100%.

Tobacco

Age of First Use

Figure 6. Age of First Use of Tobacco, 2024




Table 8. Age of First Tobacco Use, 2021-2024

Age of First Use	Year			
	2021	2022	2023	2024
Do not use/ Never	2159 (66.15%)	2038 (52.78%)	687 (60.16%)	1334 (64.32%)
Under 10	55 (1.69%)	119 (3.08%)	34 (2.98%)	39 (1.88%)
11-15	380 (11.64%)	529 (13.70%)	164 (14.36%)	241 (11.62%)
16-20	567 (17.37%)	833 (21.57%)	200 (17.51%)	366 (17.65%)
21-25	82 (2.51%)	232 (6.01%)	48 (4.20%)	81 (3.91%)
26+	21 (0.64%)	110 (2.85%)	9 (0.79%)	13 (0.63%)

Note. Due to rounding error, percentages may not sum to 100%.

Table 9. Age of First Use of Tobacco by Current Age Groups, 2024

Age of First Use	Age Groups (18-20, 21+)		Age Groups (18-25, 26+)	
	18-20	21 and Older	18-25	26 and Older
Do not use/ Never	652 (78.46%)	679 (54.89%)	1042 (73.23%)	289 (44.81%)
Under 10	9 (1.08%)	29 (2.34%)	17 (1.19%)	21 (3.26%)
11-15	62 (7.46%)	179 (14.47%)	114 (8.01%)	127 (19.69%)
16-20	108 (13.00%)	256 (20.70%)	219 (15.39%)	145 (22.48%)
21-25	N/A	81 (6.55%)	31 (2.18%)	50 (7.75%)
26+	N/A	13 (1.05%)	N/A	13 (2.02%)

Note. Due to rounding error, percentages may not sum to 100%.

Lifetime Use

Figure 7. Lifetime Occasions of Tobacco Use, 2024




Table 10. Lifetime Occasions of Use of Tobacco by Current Age Groups, 2024

Lifetime Use	Age Groups (18-20, 21+)		Age Groups (18-25, 26+)	
	18-20	21 and Older	18-25	26 and Older
0 times	658 (79.76%)	616 (50.99%)	1020 (72.75%)	254 (40.25%)

1-2 times	64 (7.76%)	111 (9.19%)	114 (8.13%)	61 (9.67%)
3-5 times	19 (2.30%)	58 (4.80%)	45 (3.21%)	32 (5.07%)
6-9 times	9 (1.09%)	39 (3.23%)	28 (2.00%)	20 (3.17%)
10+ times	75 (9.09%)	384 (31.79%)	195 (13.91%)	264 (41.84%)

Note. Due to rounding error, percentages may not sum to 100%.

Recent Use

Table 11. Tobacco Use in the Last 30 Days, 2021-2024

Recent Use	Year			
	2021	2022	2023	2024
0 times	2803 (88.26%)	2672 (70.76%)	930 (86.27%)	1766 (87.64%)
1-2 times	77 (2.42%)	236 (6.25%)	24 (2.23%)	66 (3.28%)
3-5 times	31 (0.98%)	226 (5.99%)	16 (1.48%)	28 (1.39%)
6-9 times	14 (0.44%)	163 (4.32%)	99 (9.18%)	0 (0.00%)
10+ times	251 (7.90%)	479 (12.69%)	9 (0.83%)	155 (7.69%)

Note. Due to rounding error, percentages may not sum to 100%.

Figure 8. Occasions of Tobacco Use in the Past 30 Days, 2024




Table 12. Tobacco Use in the Last 30 Days by Current Age Groups, 2024

Recent Use	Age Groups (18-20, 21+)		Age Groups (18-25, 26+)	
	18-20	21 and Older	18-25	26 or Older
0 times	750 (92.36%)	1011 (84.46%)	1251 (90.52%)	510 (81.34%)
1-2 times	20 (2.46%)	45 (3.76%)	45 (3.26%)	20 (3.19%)
3-5 times	11 (1.35%)	17 (1.42%)	22 (1.59%)	6 (0.96%)
6-9 times	0 (0.00%)	0 (0.00%)	0 (0.00%)	0 (0.00%)
10+ times	31 (3.82%)	124 (10.36%)	64 (4.63%)	91 (14.51%)

Note. Due to rounding error, percentages may not sum to 100%.

Vaping

Age of First Use

Figure 9. Age of First Use of Vape Products, 2024




Table 13. Age of First Vape Product Use, 2021-2024

Age of First Use	Year			
	2021	2022	2023	2024
Do not use/ Never	2296 (70.78%)	2349 (61.57%)	714 (62.80%)	1371 (66.72%)
Under 10	14 (0.43%)	52 (1.36%)	7 (0.62%)	11 (0.54%)
11-15	146 (4.50%)	271 (7.10%)	72 (6.33%)	146 (7.10%)
16-20	505 (15.57%)	633 (16.59%)	209 (18.38%)	341 (16.59%)
21-25	146 (4.50%)	279 (7.31%)	72 (6.33%)	89 (4.33%)
26+	137 (4.22%)	231 (6.06%)	63 (5.54%)	97 (4.72%)

Note. Due to rounding error, percentages may not sum to 100%.

Table 14. Age of First Use of Vape Products by Current Age Groups, 2024

Age of First Use	Age Groups (18-20, 21+)		Age Groups (18-25, 26+)	
	18-20	21 and Older	18-25	26 and Older
Do not use/ Never	575 (69.44%)	793 (64.95%)	921 (65.09%)	447 (70.50%)
Under 10	7 (0.85%)	4 (0.33%)	11 (0.78%)	0 (0.00%)
11-15	99 (11.96%)	46 (3.77%)	141 (9.96%)	4 (0.63%)
16-20	146 (17.63%)	194 (15.89%)	295 (20.85%)	45 (7.10%)
21-25	N/A	89 (7.29%)	46 (3.25%)	43 (6.78%)
26+	N/A	95 (7.78%)	N/A	95 (14.98%)

Note. Due to rounding error, percentages may not sum to 100%.

Lifetime Use

Figure 10. Lifetime Occasions of Using Vape Products, 2024




Table 15. Lifetime Occasions of Vape Product Use by Current Age Groups, 2024

Lifetime Use	Age Groups (18-20, 21+)		Age Groups (18-25, 26+)	
	18-20	21 and Older	18-25	26 and Older
0 times	544 (66.26%)	746 (62.95%)	870 (62.54%)	420 (68.29%)
1-2 times	58 (7.06%)	39 (3.29%)	84 (6.04%)	13 (2.11%)
3-5 times	50 (6.09%)	47 (3.97%)	78 (5.61%)	19 (3.09%)
6-9 times	18 (2.19%)	29 (2.45%)	35 (2.52%)	12 (1.95%)
10+ times	151 (18.39%)	324 (27.34%)	324 (23.29%)	151 (24.55%)

Note. Due to rounding error, percentages may not sum to 100%.

Recent Use

Table 16. Vape Product Use in the Last 30 Days, 2021-2024

Variable/Use	Year			
	2021	2022	2023	2024
Nicotine Vape				
0 times	2674 (84.35%)	2652 (70.49%)	836 (77.70%)	1629 (81.25%)
1-2 times	91 (2.87%)	195 (5.18%)	24 (2.23%)	79 (3.94%)
3-5 times	39 (1.23%)	206 (5.48%)	18 (1.67%)	26 (1.30%)
6-9 times	19 (0.60%)	156 (4.15%)	192 (17.84%)	0 (0.00%)
10+ times	347 (10.95%)	553 (14.70%)	6 (0.56%)	271 (13.52%)
Marijuana Vape				
0 times	2920 (92.35%)	2992 (79.77%)	931 (86.44%)	1773 (88.65%)
1-2 times	69 (2.18%)	179 (4.77%)	34 (3.16%)	69 (3.45%)
3-5 times	43 (1.36%)	215 (5.73%)	18 (1.67%)	25 (1.25%)
6-9 times	16 (0.51%)	141 (3.76%)	89 (8.26%)	0 (0.00%)
10+ times	114 (3.61%)	224 (5.97%)	5 (0.46%)	133 (6.65%)
Flavor Vape				
0 times	3033 (96.01%)	3288 (87.66%)	1033 (96.09%)	1898 (94.99%)
1-2 times	51 (1.61%)	196 (5.23%)	12 (1.12%)	37 (1.85%)
3-5 times	18 (0.57%)	153 (4.08%)	6 (0.56%)	19 (0.95%)
6-9 times	10 (0.32%)	53 (1.41%)	21 (1.95%)	0 (0.00%)
10+ times	47 (1.49%)	61 (1.63%)	3 (0.28%)	44 (2.20%)

Note. Due to rounding error, percentages may not sum to 100%.

Figure 11. Occasions of Vaping Nicotine in the Last 30 Days, 2024




Figure 12. Occasions of Vaping Marijuana in the Last 30 Days, 2024




Figure 13. Occasions of Vaping Flavor in the Last 30 Days, 2024




Table 17. Use of Vape for Nicotine in the Last 30 Days by Current Age Groups, 2024

Recent Use	Age Groups (18-20, 21+)		Age Groups (18-25, 26+)	
	18-20	21 and Older	18-25	26 or Older
0 times	665 (82.10%)	959 (80.66%)	1102 (80.15%)	522 (83.65%)
1-2 times	37 (4.57%)	42 (3.53%)	66 (4.80%)	13 (2.08%)
3-5 times	18 (2.22%)	8 (0.67%)	23 (1.67%)	3 (0.48%)
6-9 times	0 (0.00%)	0 (0.00%)	0 (0.00%)	0 (0.00%)
10+ times	90 (11.11%)	180 (15.14%)	184 (13.38%)	86 (13.78%)

Note. Due to rounding error, percentages may not sum to 100%.

Table 18. Use of Vape for Marijuana in the Last 30 Days by Current Age Groups, 2024

Recent Use	Age Groups (18-20, 21+)		Age Groups (18-25, 26+)	
	18-20	21 and Older	18-25	26 or Older
0 times	713 (88.57%)	1056 (88.81%)	1208 (88.11%)	561 (90.05%)
1-2 times	28 (3.48%)	40 (3.36%)	49 (3.57%)	19 (3.05%)
3-5 times	12 (1.49%)	13 (1.09%)	19 (1.39%)	6 (0.96%)
6-9 times	0 (0.00%)	0 (0.00%)	0 (0.00%)	0 (0.00%)
10+ times	52 (6.46%)	80 (6.73%)	95 (6.93%)	37 (5.94%)

Note. Due to rounding error, percentages may not sum to 100%.

Table 19. Use of Vape for Flavor in the Last 30 Days by Current Age Groups, 2024

Recent Use	Age Groups (18-20, 21+)		Age Groups (18-25, 26+)	
	18-20	21 and Older	18-25	26 or Older
0 times	766 (95.27%)	1126 (94.78%)	1299 (94.82%)	593 (95.34%)
1-2 times	15 (1.87%)	22 (1.85%)	25 (1.82%)	12 (1.93%)
3-5 times	12 (1.49%)	7 (0.59%)	18 (1.31%)	1 (0.16%)
6-9 times	0 (0.00%)	0 (0.00%)	0 (0.00%)	0 (0.00%)
10+ times	11 (1.37%)	33 (2.78%)	28 (2.04%)	16 (2.57%)

Note. Due to rounding error, percentages may not sum to 100%.

Marijuana

Age of First Use

Figure 14. Age of First Use of Marijuana, 2024




Table 20. Age of First Use of Marijuana, 2021-2024

Age of First Use	Year			
	2021	2022	2023	2024
Do not use/ Never	1999 (61.39%)	2245 (58.45%)	570 (49.96%)	1149 (55.56%)
Under 10	21 (0.64%)	60 (1.56%)	9 (0.79%)	11 (0.53%)
11-15	268 (8.23%)	415 (10.80%)	123 (10.78%)	247 (11.94%)
16-20	774 (23.77%)	827 (21.53%)	355 (31.11%)	539 (26.06%)
21-25	148 (4.55%)	206 (5.36%)	62 (5.43%)	99 (4.79%)
26+	46 (1.41%)	88 (2.29%)	22 (1.93%)	23 (1.11%)

Note. Due to rounding error, percentages may not sum to 100%.

Table 21. Age of First Use of Marijuana by Current Age Groups, 2024

Age of First Use	Age Groups (18-20, 21+)		Age Groups (18-25, 26+)	
	18-20	21 and Older	18-25	26 and Older
Do not use/ Never	557 (67.11%)	589 (47.81%)	864 (60.76%)	282 (44.06%)
Under 10	7 (0.84%)	4 (0.32%)	9 (0.63%)	2 (0.31%)
11-15	79 (9.52%)	167 (13.56%)	133 (9.35%)	113 (17.66%)
16-20	187 (22.53%)	350 (28.41%)	374 (26.30%)	163 (25.47%)
21-25	N/A	99 (8.04%)	42 (2.95%)	57 (8.91%)
26+	N/A	23 (1.87%)	N/A	23 (3.59%)

Note. Due to rounding error, percentages may not sum to 100%.

Lifetime Use

Figure 15. Lifetime Occasions of Marijuana Use, 2024




Table 22. Lifetime Occasions of Use of Marijuana by Current Age Groups, 2024

Lifetime Use	Age Groups (18-20, 21+)		Age Groups (18-25, 26+)	
	18-20	21 and Older	18-25	26 and Older
0 times	537 (65.17%)	537 (65.17%)	828 (59.19%)	244 (38.73%)
1-2 times	69 (8.37%)	69 (8.37%)	123 (8.79%)	75 (11.90%)
3-5 times	47 (5.70%)	47 (5.70%)	86 (6.15%)	40 (6.35%)
6-9 times	30 (3.64%)	30 (3.64%)	60 (4.29%)	35 (5.56%)
10+ times	141 (17.11%)	141 (17.11%)	302 (21.59%)	236 (37.46%)

Recent Use

Table 23. Marijuana Use in the Last 30 Days, 2021-2024

Recent Use	Year			
	2021	2022	2023	2024
0 times	2770 (87.52%)	3000 (79.72%)	894 (82.93%)	1687 (84.22%)
1-2 times	115 (3.63%)	277 (7.36%)	52 (4.82%)	99 (4.94%)
3-5 times	58 (1.83%)	154 (4.09%)	28 (2.60%)	34 (1.70%)
6-9 times	25 (0.79%)	91 (2.42%)	90 (8.35%)	0 (0.00%)
10+ times	197 (6.22%)	241 (6.40%)	14 (1.30%)	183 (9.14%)

Note. Due to rounding error, percentages may not sum to 100%.

Figure 16. Occasions of Marijuana Use in the Last 30 Days, 2024




Table 24. Use of Marijuana in the Last 30 Days by Current Age Groups, 2024

Recent Use	Age Groups (18-20, 21+)		Age Groups (18-25, 26+)	
	18-20	21 and Older	18-25	26 or Older
0 times	682 (84.51%)	1001 (84.12%)	1157 (84.15%)	526 (84.57%)
1-2 times	47 (5.82%)	52 (4.37%)	77 (5.60%)	22 (3.54%)
3-5 times	19 (2.35%)	15 (1.26%)	29 (2.11%)	5 (0.80%)
6-9 times	0 (0.00%)	0 (0.00%)	0 (0.00%)	0 (0.00%)
10+ times	59 (7.31%)	122 (10.25%)	112 (8.15%)	69 (11.09%)

Prescription Medication

Age of First Use

Figure 17. Age of First Use of Prescription Opioids, 2024




Figure 18. Age of First Use of Another Person's Prescription Drugs, 2024




Table 25. Age of First Use of Prescription Drugs, 2021-2024

Variable/Age	Year			
	2021	2022	2023	2024
Prescription Opioids				
Do not use/ Never	2712 (83.57%)	3038 (79.28%)	913 (80.37%)	1731 (84.32%)
Under 10	15 (0.46%)	46 (1.20%)	11 (0.97%)	7 (0.34%)
11-15	96 (2.96%)	162 (4.23%)	38 (3.35%)	65 (3.17%)
16-20	273 (8.41%)	341 (8.90%)	95 (8.36%)	162 (7.89%)
21-25	94 (2.90%)	147 (3.84%)	48 (4.23%)	57 (2.78%)
26+	55 (1.69%)	98 (2.56%)	31 (2.73%)	31 (1.51%)
Other Person's Prescriptions				
Do not use/ Never	2900 (89.53%)	3134 (82.24%)	994 (87.27%)	1848 (90.15%)
Under 10	16 (0.49%)	73 (1.92%)	12 (1.05%)	13 (0.63%)
11-15	74 (2.28%)	168 (4.41%)	36 (3.16%)	64 (3.12%)
16-20	180 (5.56%)	245 (6.43%)	62 (5.44%)	88 (4.29%)
21-25	52 (1.61%)	125 (3.28%)	27 (2.37%)	23 (1.12%)
26+	17 (0.52%)	66 (1.73%)	8 (0.70%)	14 (0.68%)

Note. Due to rounding error, percentages may not sum to 100%.

Table 26. Age of First Use of Prescription Opioids by Current Age Groups, 2024

Age of First Use	Age Groups (18-20, 21+)		Age Groups (18-25, 26+)	
	18-20	21 and Older	18-25	26 and Older
Do not use/ Never	760 (91.90%)	966 (79.18%)	1275 (90.23%)	451 (71.14%)
Under 10	4 (0.48%)	3 (0.25%)	6 (0.42%)	1 (0.16%)
11-15	22 (2.66%)	43 (3.52%)	37 (2.62%)	28 (4.42%)
16-20	41 (4.96%)	120 (9.84%)	84 (5.94%)	77 (12.15%)
21-25	N/A	57 (4.67%)	11 (0.78%)	46 (7.26%)
26+	N/A	31 (2.54%)	N/A	31 (4.89%)

Note. Due to rounding error, percentages may not sum to 100%.

Table 27. Age of First Use of Another Person's Prescription by Current Age Groups, 2024

Age of First Use	Age Groups (18-20, 21+)		Age Groups (18-25, 26+)	
	18-20	21 and Older	18-25	26 and Older
Do not use/ Never	777 (94.07%)	1065 (87.44%)	1310 (92.84%)	532 (84.04%)
Under 10	9 (1.09%)	4 (0.33%)	11 (0.78%)	2 (0.32%)
11-15	23 (2.78%)	41 (3.37%)	39 (2.76%)	25 (3.95%)
16-20	17 (2.06%)	71 (5.83%)	47 (3.33%)	41 (6.48%)

21-25	N/A	23 (1.89%)	4 (0.28%)	19 (3.00%)
26+	N/A	14 (1.15%)	N/A	14 (2.21%)

Note. Due to rounding error, percentages may not sum to 100%.

Lifetime Use

Figure 19. Lifetime Occasions of Another Person's Prescription Use, 2024




Table 28. Lifetime Occasions of Use of Another Person's Prescription by Current Age Groups, 2024

Lifetime Use	Age Groups (18-20, 21+)		Age Groups (18-25, 26+)	
	18-20	21 and Older	18-25	26 and Older
0 times	768 (93.66%)	1044 (87.66%)	1294 (92.76%)	518 (84.09%)
1-2 times	25 (3.05%)	46 (3.86%)	49 (3.51%)	22 (3.57%)
3-5 times	14 (1.71%)	36 (3.02%)	26 (1.86%)	24 (3.90%)
6-9 times	5 (0.61%)	13 (1.09%)	8 (0.57%)	10 (1.62%)
10+ times	8 (0.98%)	52 (4.37%)	18 (1.29%)	42 (6.82%)

Note. Due to rounding error, percentages may not sum to 100%.

Recent Use

Table 29. Prescription Drug and Opioid Use in the Last 30 Days, 2021-2024

Recent Use	Year			
	2021	2022	2023	2024
Another Person's Prescription				
0 times	3093 (98.16%)	3351 (89.53%)	1041 (97.38%)	1963 (98.64%)
1-2 times	35 (1.11%)	173 (4.62%)	16 (1.50%)	14 (0.70%)
3-5 times	12 (0.38%)	121 (3.23%)	7 (0.65%)	8 (0.40%)
6-9 times	3 (0.10%)	72 (1.92%)	3 (0.28%)	0 (0.00%)
10+ times	8 (0.25%)	26 (0.69%)	2 (0.19%)	5 (0.25%)
Prescription Opioids				
0 times	3090 (98.35%)	3337 (89.46%)	1040 (97.38%)	1954 (98.39%)
1-2 times	24 (0.76%)	163 (4.37%)	9 (0.84%)	14 (0.70%)
3-5 times	11 (0.35%)	112 (3.00%)	5 (0.47%)	7 (0.35%)
6-9 times	7 (0.22%)	75 (2.01%)	11 (1.03%)	0 (0.00%)
10+ times	10 (0.32%)	43 (1.15%)	3 (0.28%)	11 (0.55%)

Note. Due to rounding error, percentages may not sum to 100%.

Figure 20. Occasions of Using Another Person's Prescriptions in the Last 30 Days, 2024




Figure 21. Occasions of Prescription Opioid Use in the Last 30 Days, 2024




Table 30. Use of Prescription Opioids in the Last 30 Days by Current Age Groups, 2024

Recent Use	Age Groups (18-20, 21+)		Age Groups (18-25, 26+)	
	18-20	21 and Older	18-25	26 or Older
0 times	794 (98.76%)	1154 (98.13%)	1352 (98.90%)	596 (97.23%)
1-2 times	3 (0.37%)	11 (0.94%)	5 (0.37%)	9 (1.47%)
3-5 times	3 (0.37%)	4 (0.34%)	3 (0.22%)	4 (0.65%)
6-9 times	0 (0.00%)	0 (0.00%)	0 (0.00%)	0 (0.00%)
10+ times	4 (0.50%)	7 (0.60%)	7 (0.51%)	4 (0.65%)

Note. Due to rounding error, percentages may not sum to 100%.

Table 31. Use of Another Person's Prescription in the Last 30 Days by Current Age Groups, 2024

Recent Use	Age Groups (18-20, 21+)		Age Groups (18-25, 26+)	
	18-20	21 and Older	18-25	26 or Older
0 times	795 (98.64%)	1162 (98.64%)	1350 (98.54%)	607 (98.86%)
1-2 times	7 (0.87%)	7 (0.59%)	10 (0.73%)	4 (0.65%)
3-5 times	2 (0.25%)	6 (0.51%)	5 (0.36%)	3 (0.49%)
6-9 times	0 (0.00%)	0 (0.00%)	0 (0.00%)	0 (0.00%)
10+ times	2 (0.25%)	3 (0.25%)	5 (0.36%)	0 (0.00%)

Note. Due to rounding error, percentages may not sum to 100%.

Illegal Drugs

Age of First Use

Table 32. Age of First Use of Illegal Drugs, 2024

<u>Drug</u>	<u><i>n</i></u>	<u><i>%</i></u>
Cocaine (e.g., crack, rock, freebase)		
Do not use/ Never	1916	92.96
Under 10	1	0.05
11-15	16	0.78
16-20	67	3.25
21-25	48	2.33
26+	13	0.63
Amphetamines (e.g., diet pills)		
Do not use/ Never	1815	88.11
Under 10	16	0.78
11-15	36	1.75
16-20	94	4.56
21-25	69	3.35
26+	30	1.46
Sedatives (e.g., ludes, downers)		
Do not use/ Never	1922	93.48
Under 10	5	0.24
11-15	29	1.41
16-20	67	3.26
21-25	23	1.12
26+	10	0.49
Hallucinogens/Psychedelics (e.g., PCP, LSD)		
Do not use/ Never	1827	88.73
Under 10	3	0.15
11-15	22	1.07
16-20	143	6.95
21-25	49	2.38
26+	15	0.73
Opiates (e.g., heroin, smack)		
Do not use/ Never	2001	97.47
Under 10	1	0.05
11-15	7	0.34
16-20	28	1.36

21-25	12	0.58
26+	4	0.19
<hr/>		
Inhalants (e.g., solvents, glue, gas)		
Do not use/ Never	1975	96.06
Under 10	11	0.54
11-15	29	1.41
16-20	30	1.46
21-25	10	0.49
26+	1	0.05
<hr/>		
Designer Drugs (e.g., ecstasy, MDMA, molly)		
Do not use/ Never	1897	92.36
Under 10	2	0.10
11-15	14	0.68
16-20	84	4.09
21-25	48	2.34
26+	9	0.44
<hr/>		
Steroids		
Do not use/ Never	1996	97.22
Under 10	9	0.44
11-15	10	0.49
16-20	27	1.32
21-25	6	0.29
26+	5	0.24
<hr/>		
Methamphetamine (e.g., meth, ice, speed)		
Do not use/ Never	1970	95.82
Under 10	3	0.15
11-15	10	0.49
16-20	39	1.90
21-25	29	1.41
26+	5	0.24
<hr/>		
Other Illegal Drugs		
Do not use/ Never	1973	96.57
Under 10	3	0.15
11-15	19	0.93
16-20	33	1.62
21-25	9	0.44
26+	6	0.29

Note. Due to rounding errors, percentages may not equal 100%.

Table 33. Age of First Use of Illegal Drugs by Current Age Groups, 2024

Drug/Age of First Use	Age Groups (18-20, 21+)		Age Groups (18-25, 26+)	
	18-20	21 and Older	18-25	26 and Older
Cocaine (e.g., crack, rock, freebase)				
Do not use/ Never	818 (98.79%)	1092 (89.00%)	1377 (97.25%)	533 (83.41%)
Under 10	0 (0.00%)	1 (0.08%)	1 (0.07%)	0 (0.00%)
11-15	3 (0.36%)	13 (1.06%)	6 (0.42%)	10 (1.56%)
16-20	7 (0.85%)	60 (4.89%)	23 (1.62%)	44 (6.89%)
21-25	N/A	48 (3.91%)	9 (0.64%)	39 (6.10%)
26+	N/A	13 (1.06%)	N/A	13 (2.03%)
Amphetamines (e.g., diet pills)				
Do not use/ Never	797 (96.37%)	1012 (82.48%)	1328 (93.72%)	481 (75.51%)
Under 10	5 (0.60%)	11 (0.90%)	14 (0.99%)	2 (0.31%)
11-15	9 (1.09%)	27 (2.20%)	15 (1.06%)	21 (3.30%)
16-20	16 (1.93%)	78 (6.36%)	40 (2.82%)	54 (8.48%)
21-25	N/A	69 (5.62%)	20 (1.41%)	49 (7.69%)
26+	N/A	30 (2.44%)	N/A	30 (4.71%)
Sedatives (e.g., ludes, downers)				
Do not use/ Never	801 (97.09%)	1115 (91.02%)	1371 (96.96%)	545 (85.69%)
Under 10	4 (0.48%)	1 (0.08%)	5 (0.35%)	0 (0.00%)
11-15	10 (1.21%)	19 (1.55%)	12 (0.85%)	17 (2.67%)
16-20	10 (1.21%)	57 (4.65%)	24 (1.70%)	43 (6.76%)
21-25	N/A	23 (1.88%)	2 (0.14%)	21 (3.30%)
26+	N/A	10 (0.82%)	N/A	10 (1.57%)
Hallucinogens/Psychedelics (e.g., PCP, LSD)				
Do not use/ Never	773 (93.36%)	1050 (85.71%)	1296 (91.46%)	527 (82.86%)
Under 10	2 (0.24%)	1 (0.08%)	3 (0.21%)	0 (0.00%)
11-15	6 (0.72%)	16 (1.31%)	9 (0.64%)	13 (2.04%)
16-20	47 (5.68%)	94 (7.67%)	95 (6.70%)	46 (7.23%)

21-25	N/A	49 (4.00%)	14 (0.99%)	35 (5.50%)
26+	N/A	15 (1.22%)	N/A	15 (2.36%)
Opiates (e.g., heroin, smack)				
Do not use/ Never	822 (99.40%)	1173 (96.15%)	1399 (99.08%)	596 (93.86%)
Under 10	0 (0.00%)	1 (0.08%)	1 (0.07%)	0 (0.00%)
11-15	1 (0.12%)	6 (0.49%)	3 (0.21%)	4 (0.63%)
16-20	4 (0.48%)	24 (1.97%)	8 (0.57%)	20 (3.15%)
21-25	N/A	12 (0.98%)	1 (0.07%)	11 (1.73%)
26+	N/A	4 (0.33%)	N/A	4 (0.63%)
Inhalants (e.g., solvents, glue, gas)				
Do not use/ Never	807 (97.58%)	1162 (95.01%)	1378 (97.39%)	591 (93.07%)
Under 10	7 (0.85%)	4 (0.33%)	9 (0.64%)	2 (0.31%)
11-15	5 (0.60%)	24 (1.96%)	6 (0.42%)	23 (3.62%)
16-20	8 (0.97%)	22 (1.80%)	17 (1.20%)	13 (2.05%)
21-25	N/A	10 (0.82%)	5 (0.35%)	5 (0.79%)
26+	N/A	1 (0.08%)	N/A	1 (0.16%)
Designer Drugs (e.g., ecstasy, MDMA, molly)				
Do not use/ Never	810 (97.94%)	1082 (88.62%)	1365 (96.60%)	527 (82.99%)
Under 10	0 (0.00%)	2 (0.16%)	2 (0.14%)	0 (0.00%)
11-15	6 (0.73%)	8 (0.66%)	9 (0.64%)	5 (0.79%)
16-20	10 (1.21%)	73 (5.98%)	31 (2.19%)	52 (8.19%)
21-25	N/A	47 (3.85%)	6 (0.42%)	42 (6.61%)
26+	N/A	9 (0.74%)	N/A	9 (1.42%)
Steroids				
Do not use/ Never	806 (97.58%)	1184 (96.97%)	1382 (97.81%)	608 (95.90%)
Under 10	3 (0.36%)	6 (0.49%)	5 (0.35%)	4 (0.63%)
11-15	6 (0.73%)	4 (0.33%)	7 (0.50%)	3 (0.47%)
16-20	11 (1.33%)	16 (1.31%)	17 (1.20%)	10 (1.58%)
21-25	N/A	6 (0.49%)	2 (0.14%)	4 (0.63%)
26+	N/A	5 (0.41%)	N/A	5 (0.79%)
Methamphetamine (e.g., meth, ice, speed)				

Do not use/ Never	821 (99.39%)	1143 (93.38%)	1393 (98.58%)	571 (89.64%)
Under 10	1 (0.12%)	2 (0.16%)	2 (0.14%)	1 (0.16%)
11-15	2 (0.24%)	8 (0.65%)	4 (0.28%)	6 (0.94%)
16-20	2 (0.24%)	37 (3.02%)	13 (0.92%)	26 (4.08%)
21-25	N/A	29 (2.37%)	1 (0.07%)	28 (4.40%)
26+	N/A	5 (0.41%)	N/A	5 (0.78%)
Other Illegal Drugs				
Do not use/ Never	805 (98.17%)	1163 (95.56%)	1368 (97.57%)	600 (94.49%)
Under 10	2 (0.24%)	1 (0.08%)	3 (0.21%)	0 (0.00%)
11-15	6 (0.73%)	13 (1.07%)	11 (0.78%)	8 (1.26%)
16-20	7 (0.85%)	25 (2.05%)	18 (1.28%)	14 (2.20%)
21-25	N/A	9 (0.74%)	2 (0.14%)	7 (1.10%)
26+	N/A	6 (0.49%)	N/A	6 (0.94%)

Note. Due to rounding error, percentages may not sum to 100%.

Lifetime Use

Table 34. Lifetime Occasions of Illegal Drug Use, 2024

Drug	<i>n</i>	%
Cocaine (e.g., crack, rock, freebase)		
0 times	1872	92.26
1-2 times	34	1.68
3-5 times	27	1.33
6-9 times	20	0.99
10+ times	76	3.75
Amphetamines (e.g., diet pills)		
0 times	1801	88.63
1-2 times	37	1.82
3-5 times	23	1.13
6-9 times	24	1.18
10+ times	147	7.23
Sedatives (e.g., ludes, downers)		
0 times	1905	93.89
1-2 times	22	1.08
3-5 times	14	0.69
6-9 times	13	0.64

10+ times	75	3.70
<hr/>		
Hallucinogens/Psychedelics (e.g., PCP, LSD)		
0 times	1801	88.68
1-2 times	80	3.94
3-5 times	67	3.30
6-9 times	32	1.58
10+ times	51	2.51
<hr/>		
Opiates (e.g., heroin, smack)		
0 times	1957	96.79
1-2 times	14	0.69
3-5 times	11	0.54
6-9 times	7	0.35
10+ times	33	1.63
<hr/>		
Inhalants (e.g., solvents, glue, gas)		
0 times	1941	96.09
1-2 times	33	1.63
3-5 times	9	0.45
6-9 times	11	0.54
10+ times	26	1.29
<hr/>		
Designer Drugs (e.g., ecstasy, MDMA, molly)		
0 times	1873	92.68
1-2 times	54	2.67
3-5 times	45	2.23
6-9 times	16	0.79
10+ times	33	1.63
<hr/>		
Steroids		
0 times	1957	97.03
1-2 times	25	1.24
3-5 times	12	0.59
6-9 times	7	0.35
10+ times	16	0.79
<hr/>		
Methamphetamine (e.g., meth, ice, speed)		
0 times	1922	95.72
1-2 times	19	0.95
3-5 times	5	0.25
6-9 times	7	0.35
10+ times	55	2.74
<hr/>		
Other Illegal Drugs		

0 times	1937	96.56
1-2 times	20	1.00
3-5 times	9	0.45
6-9 times	2	0.10
10+ times	38	1.89

Note. Due to rounding errors, percentages may not equal 100%.

Table 35. Lifetime Occasions of Use of Illegal Drugs by Current Age Groups, 2024

Drug/Frequency	Age Groups (18-20, 21+)		Age Groups (18-25, 26+)	
	18-20	21 and Older	18-25	26 and Older
Cocaine (e.g., crack, rock, freebase)				
0 times	812 (98.78%)	1054 (87.76%)	1358 (97.21%)	508 (81.15%)
1-2 times	1 (0.12%)	33 (2.75%)	8 (0.57%)	26 (4.15%)
3-5 times	2 (0.24%)	25 (2.08%)	9 (0.64%)	18 (2.88%)
6-9 times	2 (0.24%)	18 (1.50%)	5 (0.36%)	15 (2.40%)
10+ times	5 (0.61%)	71 (5.91%)	17 (1.22%)	59 (9.42%)
Amphetamines (e.g., diet pills)				
0 times	793 (96.35%)	1002 (83.29%)	1311 (93.78%)	484 (77.07%)
1-2 times	4 (0.49%)	33 (2.74%)	14 (1.00%)	23 (3.66%)
3-5 times	5 (0.61%)	18 (1.50%)	11 (0.79%)	12 (1.91%)
6-9 times	3 (0.36%)	21 (1.75%)	5 (0.36%)	19 (3.03%)
10+ times	18 (2.19%)	129 (10.72%)	57 (4.08%)	90 (14.33%)
Sedatives (e.g., ludes, downers)				
0 times	802 (97.33%)	1097 (91.49%)	1355 (96.92%)	544 (87.04%)
1-2 times	4 (0.49%)	18 (1.50%)	12 (0.86%)	10 (1.60%)
3-5 times	5 (0.61%)	9 (0.75%)	5 (0.36%)	9 (1.44%)
6-9 times	5 (0.61%)	8 (0.67%)	6 (0.43%)	7 (1.12%)
10+ times	8 (0.97%)	67 (5.59%)	20 (1.43%)	55 (8.80%)
Hallucinogens/Psychedelics (e.g., PCP, LSD)				
0 times	772 (93.69%)	1025 (85.35%)	1281 (91.50%)	516 (82.56%)
1-2 times	25 (3.03%)	54 (4.50%)	47 (3.36%)	32 (5.12%)

3-5 times	14 (1.70%)	53 (4.41%)	35 (2.50%)	32 (5.12%)
6-9 times	5 (0.61%)	27 (2.25%)	16 (1.14%)	16 (2.56%)
10+ times	8 (0.97%)	42 (3.50%)	21 (1.50%)	29 (4.64%)
Opiates (e.g., heroin, smack)				
0 times	812 (99.02%)	1139 (95.23%)	1374 (98.57%)	577 (92.77%)
1-2 times	2 (0.24%)	12 (1.00%)	7 (0.50%)	7 (1.13%)
3-5 times	2 (0.24%)	9 (0.75%)	5 (0.36%)	6 (0.96%)
6-9 times	2 (0.24%)	5 (0.42%)	3 (0.22%)	4 (0.64%)
10+ times	2 (0.24%)	31 (2.59%)	5 (0.36%)	28 (4.50%)
Inhalants (e.g., solvents, glue, gas)				
0 times	800 (97.44%)	1135 (95.14%)	1360 (97.42%)	575 (93.04%)
1-2 times	9 (1.10%)	24 (2.01%)	14 (1.00%)	19 (3.07%)
3-5 times	3 (0.37%)	6 (0.50%)	6 (0.43%)	3 (0.49%)
6-9 times	3 (0.37%)	8 (0.67%)	6 (0.43%)	5 (0.81%)
10+ times	6 (0.73%)	20 (1.68%)	10 (0.72%)	16 (2.59%)
Designer Drugs (e.g., ecstasy, MDMA, molly)				
0 times	808 (98.30%)	1060 (88.85%)	1352 (96.85%)	516 (83.36%)
1-2 times	7 (0.85%)	47 (3.94%)	18 (1.29%)	36 (5.82%)
3-5 times	1 (0.12%)	43 (3.60%)	9 (0.64%)	35 (5.65%)
6-9 times	3 (0.36%)	13 (1.09%)	8 (0.57%)	8 (1.29%)
10+ times	3 (0.36%)	30 (2.51%)	9 (0.64%)	24 (3.88%)
Steroids				
0 times	798 (97.44%)	1153 (96.73%)	1361 (97.70%)	590 (95.47%)
1-2 times	11 (1.34%)	14 (1.17%)	16 (1.15%)	9 (1.46%)
3-5 times	4 (0.49%)	8 (0.67%)	6 (0.43%)	6 (0.97%)
6-9 times	1 (0.12%)	6 (0.50%)	2 (0.14%)	5 (0.81%)
10+ times	5 (0.61%)	11 (0.92%)	8 (0.57%)	8 (1.29%)
Methamphetamine (e.g., meth, ice, speed)				
0 times	814 (99.51%)	1102 (93.07%)	1367 (98.63%)	549 (89.12%)
1-2 times	0 (0.00%)	19 (1.60%)	7 (0.51%)	12 (1.95%)
3-5 times	1 (0.12%)	4 (0.34%)	1 (0.07%)	4 (0.65%)

6-9 times	0 (0.00%)	7 (0.59%)	2 (0.14%)	5 (0.81%)
10+ times	3 (0.37%)	52 (4.39%)	9 (0.65%)	46 (7.47%)
Other Illegal Drugs				
0 times	803 (98.29%)	1129 (95.44%)	1355 (97.83%)	577 (93.82%)
1-2 times	4 (0.49%)	15 (1.27%)	9 (0.65%)	10 (1.63%)
3-5 times	2 (0.24%)	7 (0.59%)	6 (0.43%)	3 (0.49%)
6-9 times	0 (0.00%)	2 (0.17%)	0 (0.00%)	2 (0.33%)
10+ times	8 (0.98%)	30 (2.54%)	15 (1.08%)	23 (3.74%)

Note. Due to rounding error, percentages may not sum to 100%.

Recent Use

Table 36. Use of Illegal Drugs in the Last 30 Days, 2024

Drug	<i>n</i>	%
Cocaine (e.g., crack, rock, freebase)		
0 times	1990	99.35
1-2 times	5	0.25
3-5 times	3	0.15
6-9 times	0	0.00
10+ times	5	0.25
Amphetamines (e.g., diet pills)		
0 times	1941	97.00
1-2 times	7	0.35
3-5 times	12	0.60
6-9 times	0	0.00
10+ times	41	2.05
Sedatives (e.g., ludes, downers)		
0 times	1977	98.85
1-2 times	7	0.35
3-5 times	3	0.15
6-9 times	0	0.00
10+ times	13	0.65
Hallucinogens/Psychedelics (e.g., PCP, LSD)		
0 times	1978	99.15
1-2 times	10	0.50
3-5 times	5	0.25
6-9 times	0	0.00

10+ times	2	0.10
<hr/>		
Opiates (e.g., heroin, smack)		
0 times	1988	99.65
1-2 times	2	0.10
3-5 times	2	0.10
6-9 times	0	0.00
10+ times	3	0.15
<hr/>		
Inhalants (e.g., solvents, glue, gas)		
0 times	1987	99.50
1-2 times	6	0.30
3-5 times	1	0.05
6-9 times	0	0.00
10+ times	3	0.15
<hr/>		
Designer Drugs (e.g., ecstasy, MDMA, molly)		
0 times	1987	99.65
1-2 times	4	0.20
3-5 times	1	0.05
6-9 times	0	0.00
10+ times	2	0.10
<hr/>		
Steroids		
0 times	1983	99.50
1-2 times	4	0.20
3-5 times	1	0.05
6-9 times	0	0.00
10+ times	5	0.25
<hr/>		
Methamphetamine (e.g., meth, ice, speed)		
0 times	1984	99.70
1-2 times	0	0.00
3-5 times	1	0.05
6-9 times	0	0.00
10+ times	5	0.25
<hr/>		
Other Illegal Drugs		
0 times	1976	99.70
1-2 times	4	0.20
3-5 times	0	0.00
6-9 times	0	0.00
10+ times	2	0.10

Note. Due to rounding errors, percentages may not equal 100%.

Table 37. Use of Illegal Drugs in the Last 30 Days by Current Age Groups, 2024

Drug/Frequency	Age Groups (18-20, 21+)		Age Groups (18-25, 26+)	
	18-20	21 and Older	18-25	26 and Older
Cocaine (e.g., crack, rock, freebase)				
0 times	804 (99.63%)	1180 (99.16%)	1366 (99.42%)	618 (99.20%)
1-2 times	1 (0.12%)	4 (0.34%)	4 (0.29%)	1 (0.16%)
3-5 times	1 (0.12%)	2 (0.17%)	2 (0.15%)	1 (0.16%)
6-9 times	0 (0.00%)	0 (0.00%)	0 (0.00%)	0 (0.00%)
10+ times	1 (0.12%)	4 (0.34%)	2 (0.15%)	3 (0.48%)
Amphetamines (e.g., diet pills)				
0 times	799 (98.89%)	1136 (95.70%)	1349 (98.04%)	586 (94.67%)
1-2 times	2 (0.25%)	5 (0.42%)	2 (0.15%)	5 (0.81%)
3-5 times	0 (0.00%)	12 (1.01%)	4 (0.29%)	8 (1.29%)
6-9 times	0 (0.00%)	0 (0.00%)	0 (0.00%)	0 (0.00%)
10+ times	7 (0.87%)	34 (2.86%)	21 (1.53%)	20 (3.23%)
Sedatives (e.g., ludes, downers)				
0 times	799 (99.01%)	1172 (98.74%)	1363 (99.20%)	608 (98.06%)
1-2 times	4 (0.50%)	3 (0.25%)	5 (0.36%)	2 (0.32%)
3-5 times	1 (0.12%)	2 (0.17%)	1 (0.07%)	2 (0.32%)
6-9 times	0 (0.00%)	0 (0.00%)	0 (0.00%)	0 (0.00%)
10+ times	3 (0.37%)	10 (0.84%)	5 (0.36%)	8 (1.29%)
Hallucinogens/Psychedelics (e.g., PCP, LSD)				
0 times	795 (98.76%)	1178 (99.49%)	1360 (99.20%)	613 (99.19%)
1-2 times	6 (0.75%)	3 (0.25%)	6 (0.44%)	3 (0.49%)
3-5 times	3 (0.37%)	2 (0.17%)	3 (0.22%)	2 (0.32%)
6-9 times	0 (0.00%)	0 (0.00%)	0 (0.00%)	0 (0.00%)
10+ times	1 (0.12%)	1 (0.08%)	2 (0.15%)	0 (0.00%)
Opiates (e.g., heroin, smack)				
0 times	804 (99.63%)	1178 (99.66%)	1366 (99.64%)	616 (99.68%)

1-2 times	1 (0.12%)	1 (0.08%)	2 (0.15%)	0 (0.00%)
3-5 times	1 (0.12%)	1 (0.08%)	1 (0.07%)	1 (0.16%)
6-9 times	0 (0.00%)	0 (0.00%)	0 (0.00%)	0 (0.00%)
10+ times	1 (0.12%)	2 (0.17%)	2 (0.15%)	1 (0.16%)
Inhalants (e.g., solvents, glue, gas)				
0 times	801 (99.38%)	1180 (99.58%)	1365 (99.42%)	616 (99.68%)
1-2 times	2 (0.25%)	4 (0.34%)	4 (0.29%)	2 (0.32%)
3-5 times	1 (0.12%)	0 (0.00%)	1 (0.07%)	0 (0.00%)
6-9 times	0 (0.00%)	0 (0.00%)	0 (0.00%)	0 (0.00%)
10+ times	2 (0.25%)	1 (0.08%)	3 (0.22%)	0 (0.00%)
Designer Drugs (e.g., ecstasy, MDMA, molly)				
0 times	806 (99.88%)	1175 (99.49%)	1371 (99.78%)	610 (99.35%)
1-2 times	0 (0.00%)	4 (0.34%)	0 (0.00%)	4 (0.65%)
3-5 times	0 (0.00%)	1 (0.08%)	1 (0.07%)	0 (0.00%)
6-9 times	0 (0.00%)	0 (0.00%)	0 (0.00%)	0 (0.00%)
10+ times	1 (0.12%)	1 (0.08%)	2 (0.15%)	0 (0.00%)
Steroids				
0 times	803 (99.63%)	1174 (99.41%)	1366 (99.56%)	611 (99.35%)
1-2 times	1 (0.12%)	3 (0.25%)	3 (0.22%)	1 (0.16%)
3-5 times	0 (0.00%)	1 (0.08%)	0 (0.00%)	1 (0.16%)
6-9 times	0 (0.00%)	0 (0.00%)	0 (0.00%)	0 (0.00%)
10+ times	2 (0.25%)	3 (0.25%)	3 (0.22%)	2 (0.33%)
Methamphetamine (e.g., meth, ice, speed)				
0 times	805 (99.88%)	1173 (99.58%)	1368 (99.85%)	610 (99.35%)
1-2 times	0 (0.00%)	0 (0.00%)	0 (0.00%)	0 (0.00%)
3-5 times	0 (0.00%)	1 (0.08%)	0 (0.00%)	1 (0.16%)
6-9 times	0 (0.00%)	0 (0.00%)	0 (0.00%)	0 (0.00%)
10+ times	1 (0.12%)	4 (0.34%)	2 (0.15%)	3 (0.49%)
Other Illegal Drugs				
0 times	800 (99.88%)	1171 (99.66%)	1360 (99.78%)	611 (99.67%)
1-2 times	0 (0.00%)	3 (0.26%)	1 (0.07%)	2 (0.33%)

3-5 times	0 (0.00%)	0 (0.00%)	0 (0.00%)	0 (0.00%)
6-9 times	0 (0.00%)	0 (0.00%)	0 (0.00%)	0 (0.00%)
10+ times	1 (0.12%)	1 (0.09%)	2 (0.15%)	0 (0.00%)

Note. Due to rounding error, percentages may not sum to 100%.

Campus Observations

Policies and Programs

Table 38. Student Awareness of Campus Policies and Programs for Substance Use, 2024

Question	<i>n</i>	%
Does your campus have alcohol, tobacco, and other drug policies?		
Yes	1561	96.78
No	52	3.22
If yes to policies, are campus policies enforced?		
Yes	1112	91.15
No	108	8.85
Does our college have an alcohol, tobacco, and drug prevention program?		
Yes	752	88.99
No	93	11.01
Do you believe your campus is concerned about the prevention of alcohol, tobacco, and drug use?		
Yes	1300	89.90
No	146	10.10
Are you actively involved in efforts to prevent alcohol, tobacco, and drug use on your campus?		
Yes	276	16.93
No	1354	83.07

Note. Due to rounding errors, percentages may not equal 100%.

Campus Environment and Student Substance Use

Table 39. Perceived Frequency of Substance Use on Campus, 2024

Substance	<i>n</i>	%
Tobacco		
Did not use	435	23.80
Once a year	95	5.20
Once/month	184	10.07
Twice/month	145	7.93
Once/week	249	13.62
3 times/ week	248	13.57
5 times/ week	118	6.46
Every day	354	19.37
Alcohol		
Did not use	252	13.83
Once a year	33	1.81
Once/month	130	7.14
Twice/month	209	11.47
Once/week	568	31.17
3 times/ week	393	21.57
5 times/ week	112	6.15
Every day	125	6.86
Marijuana		
Did not use	384	21.13
Once a year	92	5.06
Once/month	186	10.24
Twice/month	165	9.08
Once/week	339	18.66
3 times/ week	316	17.39
5 times/ week	105	5.78
Every day	230	12.66
Prescription Opioids		
Did not use	1180	65.27
Once a year	182	10.07
Once/month	155	8.57
Twice/month	84	4.65
Once/week	102	5.64
3 times/ week	47	2.60

5 times/ week	13	0.72
Every day	45	2.49
<hr/>		
Sedatives		
Did not use	1411	77.78
Once a year	133	7.33
Once/month	101	5.57
Twice/month	54	2.98
Once/week	53	2.92
3 times/ week	30	1.65
5 times/ week	10	0.55
Every day	22	1.21
<hr/>		
Cocaine		
Did not use	1340	73.75
Once a year	203	11.17
Once/month	106	5.83
Twice/month	46	2.53
Once/week	56	3.08
3 times/ week	28	1.54
5 times/ week	16	0.88
Every day	22	1.21
<hr/>		
Hallucinogens/Psychedelics		
Did not use	1243	68.60
Once a year	234	12.91
Once/month	147	8.11
Twice/month	76	4.19
Once/week	65	3.59
3 times/ week	19	1.05
5 times/ week	8	0.44
Every day	20	1.10
<hr/>		
Amphetamines		
Did not use	1371	75.58
Once a year	136	7.50
Once/month	91	5.02
Twice/month	51	2.81
Once/week	59	3.25
3 times/ week	47	2.59
5 times/ week	19	1.05
Every day	40	2.21

Inhalants		
Did not use	1410	77.90
Once a year	136	7.51
Once/month	90	4.97
Twice/month	51	2.82
Once/week	57	3.15
3 times/ week	29	1.60
5 times/ week	9	0.50
Every day	28	1.55
Methamphetamine		
Did not use	1448	80.53
Once a year	128	7.12
Once/month	84	4.67
Twice/month	37	2.06
Once/week	45	2.50
3 times/ week	29	1.61
5 times/ week	7	0.39
Every day	20	1.11
Other Illegal Drugs		
Did not use	1374	76.59
Once a year	137	7.64
Once/month	91	5.07
Twice/month	62	3.46
Once/week	54	3.01
3 times/ week	26	1.45
5 times/ week	11	0.61
Every day	39	2.17
Designer Drugs		
Did not use	1401	77.57
Once a year	165	9.14
Once/month	87	4.82
Twice/month	45	2.49
Once/week	52	2.88
3 times/ week	28	1.55
5 times/ week	6	0.33
Every day	22	1.22
Prescription Drugs		
Did not use	1222	67.78

Once a year	167	9.26
Once/month	140	7.76
Twice/month	99	5.49
Once/week	81	4.49
3 times/ week	39	2.16
5 times/ week	25	1.39
Every day	30	1.66
<hr/>		
Opiates		
Did not use	1398	77.07
Once a year	169	9.32
Once/month	91	5.02
Twice/month	50	2.76
Once/week	51	2.81
3 times/ week	28	1.54
5 times/ week	8	0.44
Every day	19	1.05
<hr/>		
Steroids		
Did not use	1337	74.40
Once a year	145	8.07
Once/month	109	6.07
Twice/month	65	3.62
Once/week	74	4.12
3 times/ week	29	1.61
5 times/ week	13	0.72
Every day	25	1.39
<hr/>		
Vaping		
Did not use	433	23.98
Once a year	27	1.50
Once/month	48	2.66
Twice/month	70	3.88
Once/week	120	6.64
3 times/ week	185	10.24
5 times/ week	170	9.41
Every day	753	41.69

Note. Due to rounding errors, percentages may not equal 100%.

Figure 22. Does the Social Atmosphere on Your Campus Promote Alcohol Use?

2024




Figure 23. Does the Social Atmosphere on Your Campus Promote Other Drug Use?

2024





Figure 24. Perception of Alcohol Use on Campus Relative to Others

2024



Camp_Comp

- Greater than other campuses
- About the same as other campuses
- Less than other campuses

Mental Health Indicators

Figure 25. Feelings of Nervousness in the Last 30 Days, 2024




Figure 26. Feelings of Hopelessness in the Last 30 Days, 2024




Figure 27. Feelings of Restlessness in the Last 30 Days, 2024




Figure 28. Feelings of Worthlessness in the Last 30 Days, 2024




Figure. 29. Feelings of Depression in the Last 30 Days, 2024





Figure 30. Feelings of Helplessness in the Last 30 Days, 2024



References:

- Agunwamba, A., Kawachi, I., Williams, D., Finney Rutten, L., Wilson, P., & Vismanath, K. (2017). Mental health, racial discrimination, and tobacco use differences across rural-urban California. *The Journal of Rural Health*, 180-189.
- Alexander, A. C., & Ward, K. D. (2018). Understanding post-disaster substance use and psychological distress using concepts from the self-medication hypothesis and social cognitive theory. *Journal of psychoactive drugs*, 50, 177-186.
- Allen, B., & Stuart, G. L. (2019). Decreased cigarette smoking but no change in use of electronic cigarettes following a university-wide smoking ban. *Substance Abuse*.
- American College Health Association. (2018). American College Health Association National College Health Assessment II: Fall 2017 Reference Group Executive Summary. Hanover, MD: American College Health Association.
- American Nonsmokers' Rights Foundation. (n.d.). *ANRF*. Retrieved from STEPS FOR ENACTING A SMOKEFREE COLLEGE CAMPUS POLICY: <https://no-smoke.org/steps-for-enacting-a-smokefree-college-campus-policy/>
- An, L., Demers, M., Kirch, M., Considine-Dunn, S., Nair, V., Dsgupta, K., . . . Ahluwalia, J. (2013). A Randomized Trial of an Avatar-Hosted Multiple Behavior Change Intervention for Young Adult Smokers. *JNCI Monographs*, 47, 209-215.
- Anderson, D. S. (2011). New Directions for substance-abuse prevention. *Change: The Magazine of Higher Learning*, 43(6), 46–55. <https://doi.org/10.1080/00091383.2011.618082>
- Anon. 2022. "JED Urges Greater Understanding of the Connection Between Substance Misuse and Youth Mental Health." The JED Foundation, October 27. <https://jedfoundation.org/news-views/jed-urges-greater-understanding-of-the-connection-between-substance-misuse-and-youth-mental-health/>
- Arria, A., Caldeira, K., Bugbee, B., Vincent, K., & O'Grady, K. (2016). Marijuana use trajectories during college predict health outcomes nine years post-matriculation. *Drug and Alcohol Dependence*, 159, 158-165.
- Arria, A. M., Caldeira, K., Bugbee, B., Vincent, K., & O'Grady, K. (2015). The academic consequences of marijuana use during college. *Psychology of Addictive Behaviors*, 29(3), 564-575.
- Arria, A., Garnier-Dykstra, L., Cook, E., Caldeira, K., Vincent, K., Baron, R., & O'Grady, K. (2013). Drug use patterns in young adulthood and post-college employment. *Drug and Alcohol Dependence*, 127(1-3), 23-30.
- Arria, A., O'Grady, K., Caldeira, K., Vincent, K., & Wish, E. (2008). Nonmedical Use of Prescription Stimulants and Analgesics: Associations with Social and Academic Behaviors among College Students. *Journal of Drug Uses*, 38(4), 1045-1060.

- Auerbach RP, Alonso J, Axinn WG, et al. Mental disorders among college students in the World Health Organization World Mental Health Surveys. *Psychological Medicine*. 2016;46(14):2955-297
- Auf, R., Trepka, M., Selim, M., Taleb, Z., De La Rosa, M., Bastida, E., & Cano, M. (2019). E-cigarette use is associated with other tobacco use among U.S. adolescents. *International Journal of Public Health*, 64, 125-134.
- Babor, T., Caetano, R., Casswell, S., Edwards, G., Giesbrecht, N., Graham, K., . . . Rossow, I. (2003). *Alcohol: No Ordinary Commodity Research and public policy, second edition*. New York: Oxford University Press.
- Barker, D. C., Schleider, N. C., Ababesh, K., Johnson, T. O., & Henriksen, L. (2018). ENDS retailers and marketing near university campuses with and without tobacco-free policies. *Tobacco Induced Diseases*, 16.
- Baskerville, N. B., Azagba, S., Norman, C., McKeown, K., & Brown, K. S. (2016). Effect of a digital social media campaign on young adult smoking cessation. *Nicotine Tobacco Research*, 18(3), 351-360.
- Bayly, J., Trad, C., Saint-Fort, L., Andrew, M., Patel, M., Haynie, D., . . . Choi, K. (2020). Adoption of electronic-cigarette-free, hookah-free and American College Health Association recommended tobacco-free policies among a national sample of postsecondary educational institutions. *Journal of American College Health*, 68(1), 26-31.
- Bennett, B. L., Deiner, M., & Pokhrel, P. (2017). College anti-smoking policies and student smoking behavior: a review of the literature. *Tob Induc Ris*, 11.
- Benson, K., Flory, K., Humphreys, K., & Lee, S. (2015, March). Misuse of stimulant medication among college students: a comprehensive review and meta-analysis. *Clin Child Fam Psychol Rev*, 18(1), 50-76.
- Berg, C. J., Sutfin, E. L., Mendel, J., & Ahluwalia, J. S. (2012). Use of and interest in smoking cessation strategies among daily and nondaily student smokers. *Journal of American College Health*, 60(3), 194-202.
- Berg, C., Stratton, E., Sokol, M., Santamaria, A., Bryant, L., & Rodriguez, R. (2014). Novel incentives and messaging in an online college smoking intervention. *American Journal of Health Behavior*, 38(5), 668-680.
- Berg, C. J., Yang, Y. T., Pratt-Chapman, M. L., Douglas Evans, W., Cupertino, A.-P., Horn, K., Bernat, D. H., Abrams, L. C., & Tercyak, K. P. (2021, April 26). Campus Tobacco Control Policies and cessation interventions in college students: A commentary calling for research and action to address tobacco-related health disparities. *Translational behavioral medicine*. Retrieved April 12, 2023, from <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC8075612/>

- Blake, k. D., Klein, A. L., Walpert, L., Casey, L., Hallett, C., Douglas, C., . . . Koh, H. K. (2018). Smoke-free and tobacco-free colleges and universities in the United States. *Tobacco Control*.
- Blanco, C., Florez-Salamanca, L., Secades-Villa, R., Wang, S., & Hasin, D. (2018). Predictors of initiation of nicotine, alcohol, cannabis, and cocaine use: Results of the National Epidemiologic Survey on Alcohol and Related Conditions (NESARC). *The American Journal on Addictions*, 27(6), 477-484.
- Borsari, B., & Carey, K. B. (2001). Peer influences on college drinking: A review of the research. *Journal of Substance Abuse*, 13(4), 391-424.
- Braverman, M. T., Geldhof, G. J., Hoogesteger, L. A., & Johnson, J. A. (2018). Predicting students' noncompliance with a smoke-free university campus policy. *Preventative Medicine*, 114, 209-216.
- Brown, J. (2013). A review of the evidence on technology-based interventions for the treatment of tobacco dependence in college health. *Worldviews Evidence Based Nursing*, 10(3), 50-62.
- Brown, P. (2016). The invisible problem? Improving students' mental health. *Higher Education Policy Institute*, 66.
- Camenga, D., Bernstein, S., Dziura, J., Fiellin, L., & Krishnan-Sarin, S. (2021). Feasibility of text messaging to augment brief advice and nicotine replacement therapy for smoking cessation in college students. *Journal of American College Health*, 69(1), 1-8.
- Carrol, D., & Cole, A. (2022). Racial/ ethnic group comparisons of quit ratios and prevalence of cessation-related factors among adults who smoke with a quit attempt. *American Journal of Drug and Alcohol Abuse*, 48(1), 58-68.
- Case, K., Crook, B., Lazard, A., & Mackert, M. (2016). Formative research to identify perceptions of e-cigarettes in college students: Implications for future health communication campaigns. *Journal of American College Health*, 64(5), 380-389.
- Center for Disease Control and Prevention National Center for Health Statistics. (2021, November 17). National Center for Health Statistics: Drug Overdose Deaths in the U.S. Top 100,000 Annually. Retrieved from CDC, National Center for Health Statistics, Office of Communication: https://www.cdc.gov/nchs/pressroom/nchs_press_releases/2021/20211117.htm
- Centers for Disease Control and Prevention. (2022, October 21). Surgeon general's advisory on e-cigarette use among youth. Centers for Disease Control and Prevention. Retrieved April 15, 2023, from https://www.cdc.gov/tobacco/basic_information/e-cigarettes/surgeon-general-advisory/index.html#:~:text=E%2Dcigarettes%20entered%20the%20U.S.,tobacco%20product%20among%20U.S.%20youth.&text=E%2Dcigarette%20use%20among%20U.S.,first%20time%20during%202015%2D2017

- Chang, R. J. (2021, June 24). Colleges ban medical marijuana use, even in states where it is legal. *Forbes*. Retrieved April 10, 2023, from <https://www.forbes.com/sites/richardjchang/2021/06/23/colleges-ban-medical-marijuana-use-even-in-states-where-it-is-legal/>
- Choi, W. S., Nazir, N., & Pancheco, C. M. (2016). Recruitment and baseline characteristics of American Indian tribal college students participating in a tribal college and behavioral survey. *Nicotine Tobacco Research, 18*(6), 1488-1493.
- Clements, R. (1999). Prevalence of alcohol-use disorders and alcohol-related problems in a college student sample. *Journal of American College Health, 111-118*.
- Cleveland, M. J., Turrisi, R., Reavy, R., Ackerman, S., & Buxton, O. M. (2018). Examining Parent and Peer Influences of Alcohol Use: A Comparison of First-Year Community College and Baccalaureate Students. *Journal of Alcohol and Drug Education, 62*(2), 64–89. <https://www.jstor.org/stable/48511452>
- Coday, M., Richey, P., & Thomas, F. (2016). The recruitment experience of a randomized clinical trial to aid young adult smokers to stop smoking without weight gain interactive technology. *Contemporary Clinical Trials Communication, 2*, 61-68.
- Coleman, B. N., Apelberg, B. J., Ambrose, B. K., Green, K. M., Choiniere, C. J., & Bunnell, R. (2015). Association between electronic cigarette use and openness to cigarette smoking among US young adults. *Nicotine and Tobacco Research, 17*(2), 212-218.
- Copeland, A. L., Peltier, M. R., & Waldo, K. (2017). Perceived risk and benefits of e-cigarette use among college students. *Addiction and Behaviors, 71*, 31-37.
- Cremins, J., Usdan, S., Umstattd, M. R., Talbott, L. L., Turner, L. W., & Perko, M. A. (2011). Challenges and Recommendations to Enforcement of Alcohol Policies on College Campuses: An Administrator's Perspective. *Journal of American College Health, 59*(5), 427–430. <https://doi.org/10.1080/07448481.2010.502201>
- Crosnoe, R., Kendig, S., & Benner, A. (2017). College-going and trajectories of drinking from adolescence into adulthood. *Journal of Health and Social Behavior, 58*(2), 252–269. <https://doi.org/10.1177/0022146517693050>
- Cullen, K., Gentzke, A., Sawdey, M., Chang, J., Anic, G., Wang, T., . . . King, B. (2019). e-Cigarette Use Among Youth in the United States, 2019. *JAMA, 322*(21), 2095-2103.
- Cunningham, T., Croft, J., Liu, Y., Lu, H., Eke, P., & Giles, W. (2017). Racial disparities in age-specific mortality among blacks or African Americans - United States, 1999-2015. *Vital Signs. MMWR Morbidity and Mortality Weekly Recap, 66*(17), 444-456.
- Cvetovski, S., Reavley, N., & Jorm, A. (2012). The prevalence and correlates of psychological distress in Australian tertiary students compared to their community peers. *Australian & New Zealand Journal of Psychiatry, 46*, 457-467.

- Dai, H., & Hao, J. (2017). Geographic density and proximity of vape shops to colleges in the USA. *Tobacco Control, 26*, 379-285.
- Dai, H., Ramos, A., Faseru, B., Hill, J., & Sussman, S. (2021). Racial disparities of e-cigarette use among US youths: 2014-2019. *American Journal of Public Health, 2050-2058*.
- Derefinko, K. J., Bursac, Z., Mejia, M. G., Milich, R., & Lyman, D. R. (2018). Rural and urban substance use differences: effects of the transition to college. *American Journal Drug and Alcohol Abuse, 44(2)*, 224-234.
- DeJong, W., & Langford, L. (2002). A typology for campus-based alcohol prevention: Moving toward environmental management strategies. *Journal of Studies on Alcohol aSupplement, 14*, 140-147.
- DiMaggio, C., Galea, S., & Li, G. (2009). Substance use and misuse in the aftermath of terrorism. A Bayesian meta-analysis. *Addiction, 104*, 894-904.
- Donaldson, C., Stuppelbeen, D., Fecho, C., Ta, T., Zhang, X., & Williams, R. (2022). Nicotine vaping for relaxation and coping: Race/ ethnicity differences and social connectedness mechanisms. *Addictive behaviors*.
- Drug-Free Schools and Communities Act Amendments of 1989, H.R. 3614 - 101st Congress (1989). Retrieved April 14, 2023, from Text - H.R.3614 - 101st Congress (1989-1990): Drug-Free Schools and Communities Act Amendments of 1989 | Congress.gov | Library of Congress
- Eisenberg, D., Gollust, S., Golberstein, E., & Hefner, J. (2007). Prevalence and correlates of depression, anxiety, and suicidality among university students. *American Journal of Orthopsychiatry*.
- Faden, V., & Baskin, M. (2002). An evaluation of college online alcohol policy information. *Journal of American College Health, 12(2)*, 108-117.
- Fahey, M., Morris, J., Robinson, L., & Pebley, K. (2021). Association between perceived discrimination and vaping among college students. *Substance Use and Misuse, 738-741*.
- Fallin, A., Murray, M., Johnson, A. O., Riker, C. A., Rayens, M. K., & Hahn, E. J. (2012). Measuring compliance with tobacco-free campus policy. *Journal of American College Health, 60(7)*, 496-504.
- Federal Drug Administration. (2022, June 29). E-Cigarettes, Vapes, and other Electronic Nicotine Delivery Systems (ENDS). Retrieved April 16, 2023, from <https://www.fda.gov/tobacco-products/products-ingredients-components/e-cigarettes-vapes-and-other-electronic-nicotine-delivery-systems-ends>
- Federal Drug Administration. (2022, December 20). Results from the annual National Youth Tobacco Survey (NYTS). U.S. Food and Drug Administration. Retrieved April 15, 2023,

from <https://www.fda.gov/tobacco-products/youth-and-tobacco/results-annual-national-youth-tobacco-survey>

- Franks, A., Hawes, W., McCain, K., & Payakachat, N. (2017). Electronic cigarette use, knowledge, and perceptions among health professional students. *Current in Pharmacy Teaching and Learning*, 9(6), 1003-1009.
- Fromme, K., Corbin, W. R., & Kruse, M. I. (2008).
- Firkey, M. K., Sheinfil, A. Z., & Woolf-King, S. E. (2021). Substance use, sexual behavior, and general well-being of U.S. college students during the COVID-19 pandemic: A brief report. *Journal of American College Health*.
- Franks, A., Hawes, W., McCain, K., & Payakachat, N. (2017). Electronic cigarette use, knowledge, and perceptions among health professional students. *Current in Pharmacy Teaching and Learning*, 9(6), 1003-1009.
- Gledhill-Hoyt, J., Lee, J., Strote, J., & Wechsler, H. (2000). Increased use of marijuana and other illicit drugs at US colleges in the 1990s: Results of three national surveys. *Addiction*, 1655-1667.
- Gulliver, A., Farrer, L., Chan, J. K., Tait, R., Bennett, K., Calcar, A., & Griffiths, K. (2015). Technology-based interventions for tobacco and other drug use in university and college students: a systematic review and meta-analysis. *Addiction Science and Clinical Practice*, 5.
- Gunn, R. L., Sokolovsky, A. W., Drohan, M. M., Boyle, H. K., Stevens, A. K., White, H. R., & Jackson, K. (2022). The role of alcohol and cannabis co-use in drinking rate and its impact on consequences. *Alcoholism, clinical and experimental research*, 46(11), 2110–2120. Retrieved April 16, 2023, from <https://doi.org/10.1111/acer.14946>
- Hadland, S., Wharam, J., Schuster, M., Zhang, F., Samet, J., & Larochelle, M. (2017). Trends in Receipt of Buprenorphine and Naltrexone for Opioid Use Disorder Among Adolescents and Young Adults, 2001-2014. *JAMA Pediatrics*, 171(8), 747-755.
- Hallock, R., Dean, A., Knecht, Z., Spencer, J., & Taverna, E. (2013, June). A survey of hallucinogenic mushroom use, factors related to use, and perceptions of use among college students. *Drug and Alcohol Dependence*, 130(1-3), 245-248.
- Hammond, D., Reid, J. L., Rynard, V. L., Fong, G. T., Cummings, K. M., McNeill, A., Hitchman, S., Thrasher, J. F., Goniewicz, M. L., Bansal-Travers, M., O'Connor, R., Levy, D., Borland, R., & White, C. M. (2019). Prevalence of vaping and smoking among adolescents in Canada, England, and the United States: repeat national cross-sectional surveys. *BMJ (Clinical research ed.)*, 365, 12219. Retrieved April 16, 2023, from <https://doi.org/10.1136/bmj.12219>
- Hawkins, A. (1989, November 15). H.R.3614 - 101st Congress (1989-1990): Drug-Free Schools and ... H.R.3614 - Drug-Free Schools and Communities Act Amendments of 1989.

Retrieved April 10, 2023, from <https://www.congress.gov/bill/101st-congress/house-bill/3614/summary/35>

- Harries, M., Lust, K., Christenson, G., Redden, S., & Grant, J. (2018). Prescription opioid medication misuse among university students. *American Journal of Addictions*, 27(8), 618-624.
- Hawkins, J., Catalano, R., & Miller, J. (1992). Risk and protective factors for alcohol and other drug problems in adolescence and early adulthood: Implications for substance abuse prevention. *Psychological Bulletin*, 112(1), 64-105.
- Hefner, K. R., Solazzo, A., Mullaney, S., Coker, K. L., & Sofuoglu, M. (2019). E-cigarettes, alcohol use, and mental health: Use of perceptions of e-cigarettes among college students, by alcohol use and mental health status. *Addictive Behaviors*, 12-20.
- Hingson, R., Zha, W., & Weitzman, E. (2009). Magnitude of and trends in alcohol-related mortality and morbidity among U.S. college students ages 18–24, 1998–2005. *Journal of Studies on Alcohol and Drugs*, 16, 12-20.
- Hooper, M., Carpenter, K., Salmon, E., & Resnicow, K. (2023). Enhancing tobacco quitline outcomes for African American adults: An RCT of a culturally specific intervention. *American Journal of Preventive Medicine*, 65(6), 964-972.
- Hopkins, G. L., Freier, M. C., Babikian, T., Helm, H. W., McBride, D. C., Boward, M., Gillespie, S., & DiClemente, R. (2004). Substance use among students attending a Christian University that strictly prohibits the use of substances. *Journal of Research on Christian Education*, 13(1), 23–39. <https://doi.org/10.1080/10656210409484958>
- Horsman, E. N. (2014). Mental health impairment and human capital acquisition: Underage drinking as a predictor of concomitant alcohol dependence and poor educational attainment [Unpublished doctoral dissertation]. Southern Illinois University-Carbondale, Illinois. <https://opensiuc.lib.siu.edu/dissertations/923/>
- Hysenbegasi, A., Hass, S., & Rowland, C. (2005). The impact of depression on the academic productivity of university students. *Journal of Mental Health Policy Economics*, 8, 145-151.
- Ibrahim, A., Kelly, S., Adams, C., & Glazebrook, C. (2013). A systematic review of studies of depression prevalence in university students. *Journal of Psychiatry Research*, 391-400.
- Ickes, M. J., Wiggins, A. T., Rayens, M. K., & Hahn, E. J. (2020). Student Tobacco Use Behaviors on College Campuses by Strength of Tobacco Campus Policies. *American Journal of Health Promotion*, 34(7).
- Jeffries, E., Lemke, A., Shah, S., Dean, K., Richter, A., & Buckner, J. (2016). Addictive Behavior Interventions Among College Students. *Addiction and College Students*, 368-377.

- Jernigan, D. H., Shields, K., Mitchell, M., & Arria, A. M. (2019). Assessing Campus Alcohol Policies: Measuring Accessibility, Clarity, and Effectiveness. *Alcoholism, Clinical and Experimental Research*, 43(5), 1007–1015. <https://doi.org/10.1111/acer.14017>
- Johnson, S. (2019). Cannabis on campus? Not allowed. Cannabis on Campus? Not Allowed | Campus Drug Prevention. Retrieved April 10, 2023, from <http://www.campusdrugprevention.gov/content/cannabis-campus-not-allowed>
- Johnstad, P. (2018). Powerful substances in tiny amounts: An interview study of psychedelic microdosing. *Nordic Studies on Alcohol and Drugs*, 35(1), 39-51.
- Johnston, L., O'Malley, P., & Bachman, J. (2011). *Monitoring the future: National results on adolescent drug use: Overview of key findings, 2010*. Ann Arbor: Institute for Social Research, The University of Michigan.
- Johnston, L., O'Malley, P., Miech, R., Bachman, J., & Schulenberg, J. (2016). Monitoring the Future National Survey Results on Drug Use, 1975-2015: Overview, Key Findings on Adolescent Drug Use. Ann Arbor: Institute for Social Research, The University of Michigan.
- Jones, C., Hill, M., Pardini, D., & Meier, M. (2016). Prevalence and correlates of vaping cannabis in a sample of young adults. *Psychology of Addictive Behaviors*, 30(8), 915-921.
- Jones, R. D., Asare, M., & Lanning, B. (2020). A retrospective cross-sectional study on the prevalence of e-cigarette use among college students. *Journal of Community Health*, 46(1), 195–202. <https://doi.org/10.1007/s10900-020-00869-x>
- Jun, Kasperski, S., Vincent, K., Caldeira, K., Garnier-Dykstra, L., O'Grady, K., & Arria, A. (2011). College students' use of cocaine: Results from a longitudinal study. *Addictive Behaviors*, 36(4), 408-411.
- Jun, J., & Kim, J. (2020). How do colleges communicate about E-cigarettes? The presentation of risk, policy, and cessation resources on college websites. *Journal of American College Health*, 1-8.
- Kasimanickam, M., Kerr, D., Killion, R., Naimi, T., Lira, M., & Bae, H. (2024). State alcohol policy environment of U.S. colleges: Predictors of sexual assault and alcohol-related arrest and disciplinary action. *American Journal fo Preventive Medicine*, 66(1), 1-9.
- Kapner, D. A. (2008). Alcohol and other drug use at Historically Black Colleges and Universities. *PsycEXTRA Dataset*, 1–4. <https://doi.org/10.1037/e559152009-001>
- Karekla, M., & Savvides, S. (2019). Smoking cessation avatar-led Acceptance and Commitment Therapy digital intervention: feasibility and acceptability in young adults. *Translational Behavior Medicine*.

- Kates, F. R., Salloum, R. G., Thrasher, J. F., Islam, F., Fleischer, N. L., & Maziak, W. (2016). Geographic proximity of waterpipe smoking establishments to colleges in the US. *American Journal of Preventative Medicine*, 50(1), e9-e14.
- Kathol, N., & Sgoutas-Emch, S. (2016). Alcohol use in college: The relationship between religion, spirituality, and proscriptive attitudes toward alcohol. *Journal of Religion and Health*, 56(2), 437–449. <https://doi.org/10.1007/s10943-016-0210-2>
- Kava, C. M., Soule, E. K., Seegmiller, L., Gold, E., Snipes, W., Westfield, T., Wick, N., & Afifi, R. (2021). "Taking Up a New Problem": Context and Determinants of Pod-Mod Electronic Cigarette Use Among College Students. *Qualitative Health Research*, 31(4), 703–712. Retrieved April 16, 2023, from <https://doi.org/10.1177/1049732320971236>
- Kessler, R., Anthony, J., De Graaf, R., Demyttenaere, K., Gasque, I., & Unstun, T. (2007). Lifetime prevalence and age-of-onset distribution of mental disorders in the WORLD HEALTH ORGANIZATION. *World Psychiatry* 2007, 168-176.
- Kremer, M., & Levy, D. (2008). Peer effects and alcohol use among college students. *Journal of Economic Perspectives*, 22(3), 189–206. <https://doi.org/10.1257/jep.22.3.189>
- Kulak, J., Cornelius, M., Fong, G., & Giovino, G. (2016). Differences in quit attempts and cigarette smoking abstinence between whites and African Americans in the United States: literature review and results from the international tobacco control US survey. *Nicotine Tobacco Research*, S79-S87.
- Lanza, H. I., & Teeter, H. (2018). Electronic nicotine delivery systems (e-cigarette/ vape) use and co-occurring health-risk behaviors among an ethnically diverse sample of young adults. *Substance Use and Misuse*, 53(1), 154-161.
- Leavens, E. L., Lechner, W. V., Stevens, E. M., Miller, M. B., Meier, E., Brett, E. I., . . . Wagener, T. L. (2020). Electronic cigarette and combustible cigarette use following a campus-wide ban: Prevalence of use and harm perceptions. *Journal of American College Health*, 68(4), 332-335.
- Lee, Y., & Lee, K.-S. (2019). Association of depression and suicidality with electronic and conventional cigarette use in South Korean adolescents. *Substance Use and Misuse*, 54(6), 934-943.
- Lee, J., Kang, H., Bell, R., & Marmot, M. (2014). Social determinants of mental health. *International Review of Psychiatry*, 26(4), 392-407.
- Lenk, K., Erickson, D., Nelson, T., Winter, K., & Toomey, T. (2012). Alcohol policies and practices among four-year colleges in the United States: Prevalence and patterns. *Journal of Studies on Alcohol and Drugs*, 361-367.
- Lenk, K., Rode, P., Fabian, L., Bernat, D., Klein, E., & Forster, J. (2012). Cigarette use among young adults: comparisons between 2-year versus 4-year college students. *Journal of American College Health*, 60(4), 303-308.

- Leventhal, A., Dai, H., & Higgins, S. (2022). Smoking cessation prevalence and inequities in the United States. *Journal National Cancer Institute, 114*(3), 381-390.
- Levy, K., O'Grady, K., Wish, E., & Arria, A. (2005). An In-Depth Qualitative Examination of the Ecstasy Experience: Results of a Focus Group with Ecstasy-Using College Students. *Substance Use and Misuse, 40*, 9-10.
- Lim, G., Tam, W., Lu, Y., Ho, C., Zhang, M., & Ho, R. (2018). Prevalence of depression in the community from 30 countries between 1994 and 2014. *Scientific Reports, 8*, 2861.
- Lipperman-Kreda, S., Grube, J., & Paschall, M. (2010). Community Norms, Enforcement of Minimum Legal Drinking Age Laws, Personal Beliefs and Underage Drinking: An Explanatory Model. *Community Health, 35*(3), 249-257.
- Loukas, A., Batanova, M., Fernandez, A., & Agarwal, D. (2015, October). Changes in use of cigarettes and non-cigarette alternative products among college students. *Addictive Behaviors, 49*, 46-51.
- Luo, C., Zheng, X., Zeng, D. D., & Leischow, S. (2014). Portrayal of electronic cigarettes on YouTube. *BMC Public Health, 14*(1), 1028.
- Lupton, J. R., & Townsend, J. L. (2015). A systematic review and meta-analysis of the acceptability and effectiveness of university smoke-free policies. *Journal of American College Health, 63*(4), 238-247.
- Marshall, B., Toberts, K., Donnelly, J., & Rutledge, I. (2011). College student perceptions on campus alcohol policies and consumption patterns. *Journal of Drug Education, 41*(4), 345-358.
- Marshall, L., Zhang, L., Malarcher, A., Mann, N., King, B., & Alexander, R. (2017). Race/ ethnic variations in quitline use among US adult tobacco users in 45 states, 2011-2013. *Nicotine Tobacco Research, 19*(12), 1473-1481.
- Mason, M., Ola, B., Zaharakis, N., & Zhang, J. (2015). Text messaging interventions for adolescent and young adult substance use: a meta-analysis. *Prevention Science, 16*(2), 181-188.
- McIntosh, S., Wall, A. F., & Johnson, T. (2016). Tobacco control at community colleges: context and opportunities. *Tobacco Prevention and Cessation, 2*.
- Megivern, D., Pellerito, S., & Mowbray, C. (2003). Barriers to higher education for individuals with psychiatric disabilities. *Psychiatry Rehabilitation Journal, 26*, 217-231.
- Miech, Patrick, M. E., O'Malley, P. M., Johnston, L. D., & Bachman, J. G. (2020). Trends in Reported Marijuana Vaping Among U.S. Adolescents, 2017-2019. *JAMA: The Journal of the American Medical Association, 323*(5), 475-476. <https://doi.org/10.1001/jama.2019.2018>

- Miech, R., Johnston, L., O'Malley, P., Bachman, J., & Patrick, M. (2019, October 10). Trends in Adolescent Vaping, 2017–2019. *New England Journal of Medicine*, 381, 1490-1491.
- Miech, R., Patrick, M., O'Malley, P., & Johnston, L. (2017). What are kids vaping? Results from a national survey of U.S. adolescents. *Tobacco Control*, 26, 386-391.
- Mitchell, R., Toomey, T., & Erickson, D. (2005). Alcohol policies on college campuses. *Journal of American College Health*, 53(4), 149-157.
- National Institute of Alcohol Abuse and Alcoholism. (2002). *A call to action: Changing the culture of drinking at U.S. colleges*. NIH Publication 02-5010. US Department of Health and Human Services.
- National Institute on Drug Abuse of USA. (2021, 11 30). COVID-19 resources. Retrieved from National Institute on Drug Abuse of USA: <https://www.drugabuse.gov/related-topics/covid-19-resources>
- Navon, L., Jones, C. M., Ghinai, I., King, B. A., Briss, P. A., Hacker, K. A., & Layden, J. E. (2019). Risk Factors for E-Cigarette, or Vaping, Product Use-Associated Lung Injury (EVALI) Among Adults Who Use E-Cigarette, or Vaping, Products - Illinois, July-October 2019. *MMWR. Morbidity and mortality weekly report*, 68(45), 1034–1039. Retrieved April 16, 2023, from <https://doi.org/10.15585/mmwr.mm6845e1>
- Neighbors, C., Walters, S., Lee, C., Vader, A., Vehige, T., Szigethy, T., & DeJong, W. (2007). Event-specific prevention: Addressing college student drinking during known windows of risk. *Addictive Behaviors*, 32(11), 2667-2680.
- Nierenberg, A. (2018, October 8). Students' pot use puts colleges' funding at risk - The Boston Globe. *BostonGlobe.com*. Retrieved April 10, 2023, from: <https://www.bostonglobe.com/metro/2018/10/07/students-pot-use-puts-colleges-funding-risk/IDEMv1qY4LwZUsjhBDWVJK/story.html>
- Office of the Surgeon General. (n.d) Surgeon General's Advisory on E-cigarette Use Among Youth. Retrieved April 16, 2023, from: <https://e-cigarettes.surgeongeneral.gov/documents/surgeon-generals-advisory-on-e-cigarette-use-among-youth-2018.pdf>
- O'Grady, K., Arria, A., Fitzelle, D., & Wish, E. (2008). Heavy Drinking and Polydrug Use among College Students. *Journal of Drug Issues*, 38(2), 445-465.
- Oh, K., Lee, C., Oh, B., Oh, S.-W., Joh, H.-K., Choi, H., . . . Lea, S. (2019). The Relationship between Electronic Cigarette Use with or without Cigarette Smoking and Alcohol Use among Adolescents: Finding from the 11th Korea Youth Risk Behavior Web-based Survey. *Korean Journal of Family Medicine*, 40(4), 241-247.
- Omoike, O., & Johnson, K. R. (2020). Prevalence of Vaping and Behavioral Associations of Vaping Among a Community of College Students in the United States. *Journal of Community Health*, 190-194.

- Perea, S. D. (2018). Police Chief Magazine|Topics|Drugs & Alcohol|Marijuana, Law Enforcement, Mental Health: A Dangerous Social Experiment Share Marijuana, Law Enforcement, Mental Health: A Dangerous Social Experiment. <http://www.policechiefmagazine.org/marijuana-law-enforcement-mental-health/>
- Popp, J., Nyman, J. A., Luo, X., Bengtson, J., Lust, K., An, L., . . . Thomas, J. L. (2018). Cost-effectiveness of enhancing a Quit-and-Win smoking cessation program for college students. *The European Journal of Health Economics*, *19*, 1319-1333.
- Qiu, J., Shen, B., Zhao, M., Wang, Z., Xie, B., & Xu, Y. (2020). A nationwide survey of psychological distress among Chinese people in the COVID-19 epidemic: implications and policy recommendations. *General Psychiatry*, *33*(2), e100213.
- Rabius, V., Wiatrek, D., & McAlister, A. (2012). African American participation and success in telephone counseling for smoking cessation. *Nicotine Tobacco Research*, *240-242*.
- Ramo, D. E., Thrul, J., Delucchi, K. L., Hall, S., Ling, P. M., Belohlavek, A., & Prochaska, J. J. (2018). A randomized controlled evaluation of the tobacco status project, a Facebook intervention for young adults. *Addiction*.
- Rimsza, M., & Moses, K. (2005). Substance Abuse on the College Campus. *Pediatric Clinics*, *52*(1), 307-319.
- Ringwalt, C., Paschall, M., & Gitelman, A. (2011). Alcohol prevention strategies on college campuses and student alcohol abuse and related problems. *Journal of Drug Education*, *41*(1), 99-118.
- Roberts, M. E., Colby, S. M., Lu, B., & Ferketick, A. K. (2016). Understanding tobacco use onset among African Americans. *Nicotine Tobacco Research*, *S49-S56*.
- Rodgers, K. C. (2012). A review of multicomponent interventions to prevent and control tobacco use among college students. *Journal of American College Health*, *60*(3), 257-261.
- Roditis, M. L., Wang, D., Glantz, S. A., & Fallin, A. (2015). Evaluating California campus tobacco policies using the American College Health Association guidelines and the Institutional Grammar Tool. *Journal of American College Health*, *63*(1), 57-67.
- Russette, H. C., Harris, K. J., Schuldberg, D., & Green, L. (2014). Policy compliance of smokers on a tobacco-free university campus. *Journal of American College Health*, *62*(2), 110-116.
- Sanders, A., Stogner, J., & Miller, B. (2013). Perception vs. Reality: An Investigation of the Misperceptions concerning the Extent of Peer Novel Drug Use. *Journal of Drug Education*, *43*(2), 97-120.
- Schoenborn, C., & Gindi, R. (2016). Cigarette smoking status among current adult E-cigarette users, by age group - national health interview survey, united states, 2015. *MMWR-Morbidity and Mortality Weekly Report*, *65*(42), 1177.

- Schulenberg, J., Johnston, L., O'Malley, P. M., & Bachman, J. (2017). Monitoring the Future National Survey Results on Drug Use, 1975–2016: 2017 vol. II. College Students and Adults Ages 19–55. Retrieved 11 2021, from Ann Arbor, MI, University of Michigan, Institute for Social Research: http://www.monitoringthefuture.org/pubs/monographs/mtf-vol2_2017.pdf
- Schulenberg, J., Patrick, M., Johnston, L., O'Malley, P., Bachman, J., & Miech, R. (2020). Monitoring the Future national survey results on drug use, 1975–2020: Volume II, College students and adults ages 19–60. Ann Arbor: Institute for Social Research, The University of Michigan. Retrieved from http://www.monitoringthefuture.org/pubs/monographs/mtf-vol2_2020.pdf
- Sheffield, F., Darkes, J., Del Boca, F., & Goldman, M. (2010). Binge Drinking and Alcohol-Related Problems Among Community College Students: Implications for Prevention Policy. *Journal of American College Health*, 137-141.
- Sheldon, E., Simmonds-Buckley, M., Bone, C., Mascarenhas, T., Chan, N., Wincott, M., . . . Barkham, M. (2021). Prevalence and risk factors for mental health problems in university undergraduate students: A systematic review with meta-analysis. *Journal of Affective Disorders*, 287, 282-292.
- Simmons, V., Heckman, B., Fink, A., Small, B., & Brandon, T. (2013). Efficacy of an experiential, dissonance-based smoking intervention for college students delivered via the internet. *Journal of Consulting and Clinical Psychology*, 81(5), 810-820.
- Skidmore, C., Kaufman, E., & Crowell, S. (2016). Substance Use Among College Students. *Child and Adolescent Psychiatric Clinics of North America*, 25(4), 735-753.
- Sood, A., Andoh, J., Verhulst, S., Ganesh, M., Edson, B., & Hopkins-Price, P. (2009). Real world effectiveness of reactive telephone counseling for smoking cessation: a randomized controlled trial. *Chest*, 1229-1236.
- Suls, J. M., Luger, T. M., Curry, S. J., Mermelstein, R. J., Sporer, A. K., & An, L. C. (2012). Efficacy of smoking-cessation interventions for young adults: a meta analysis. *American Journal of Preventative Medicine*, 42(6), 655-662.
- Sun, Y., Bao, Y., & Kosten, T. (2020). Challenges to opioid use disorders during COVID-19. *American journal on addictions*, 29, 174-175.
- Sun, Y., Li, Y., Bao, Y., Meng, S., Sun, Y., Schumann, G., . . . Shi, J. (2020). Brief Report: Increased Addictive Internet and Substance Use Behavior During the COVID-19 Pandemic in China. *The American Journal of Addictions*, 29(4), 268-270.
- Sutfin, E. L., McNamara, R. S., Blocker, J. N., Ip, E. H., O'Brien, M. C., & Wolfson, M. C. (2012). Screening and brief intervention for tobacco use by student health providers on college campuses. *Journal of American College Health*, 60(1), 66-73.

- Sutfin, E., Swords, D., Song, E.-Y., Reboussin, B., Helme, D., Klein, E., & Wolfson, M. (2015). Screening and Counseling for Tobacco Use in Student Health Clinics: Reports of Health Care Providers. *American Journal of Health Promotion, 30*(1), e41-e49.
- Tavollaci, M.-P., Vasilio, A., Romo, L., Kotbagi, G., Kern, L., & Ladner, J. (2016). Patterns of electronic cigarette use in current and ever users among college students in France: a cross-sectional study. *BMJ Open 2016*, e011344.
- Thomas, J. L., Luo, X., Bengston, J. E., Weber-Main, A. M., Lust, K., Ahluwalia, J., & An, L. (2016). “Quit & Win” Contests Among College Students: Predictors of Long-Term Smoking Abstinence. *Journal of Health Promotion, 30*(4), 264-271.
- Thomas, J. L., Luo, X., Bengtson, J., Wang, Q., Ghidei, W., Nyman, J., . . . Ahluwalia, J. S. (2016). Enhancing Quit & Win contests to improve cessation among college smokers: a randomized clinical trial. *Addiction, 111*(2), 331-339.
- Toomey, T., & Wagenaar, A. (2002). Environmental policies to reduce college drinking: options and research findings. *Journal of Student Alcohol Supplement, 14*, 193-205.
- Toomey, T., Miazga, M., Lenk, K., Erickson, D., & Winters, K. (2011). Enforcing alcohol policies on college campuses: Reports from college enforcement officials. *Journal of Drug Education, 41*(3), 327-344.
- Toomey, T. L., Lenk, K. M., & Wagenaar, A. C. (2007). Environmental policies to reduce college drinking: an update of research findings. *Journal of studies on alcohol and drugs, 68*(2), 208–219. Retrieved April 16, 2023, from <https://doi.org/10.15288/jsad.2007.68.208>
- Trad, C., Bayly, J., Saint-Fors, L., Andrews, M., Patel, M., Sabado-Liwag, M., . . . Choi, K. (2018). Adoption of Tobacco- and Smoke-Free Policies in a US National Sample of Postsecondary Educational Institutions. *American Journal of Public Health, 108*(10), 1366-1369.
- Trumbo, C. (2017). Influence of Risk Perception on Attitudes and Norms Regarding Electronic Cigarettes. *Risk Analysis, 38*(5), 906-916.
- Turrisi, R., Padilla, A., & Wiersma, K. (2000). College student drinking: an examination of theoretical models of drinking tendencies in freshmen and upperclassmen. *Journal of Studies on Alcohol and Drugs, 61*(4), 598-602.
- U.S. Department of Health and Human Services. (2016). *E-cigarette use among youth and young adults. A report of the Surgeon General*. Atlanta, GA: US Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion.
- U.S. Department of Health & Human Services, National Institutes of Health. (2021, September 8). Marijuana use at historic high among college-aged adults in 2020. Retrieved from National Institute on Drug Abuse (NIDA) News Releases: <https://www.nih.gov/news-events/news-releases/marijuana-use-historic-high-among-college-aged-adults-2020>

- U.S. Department of Health and Human Services, Substance Abuse and Mental Health Services Administration. (2017). Results from the 2017 National Survey on Drug Use and Health: Detailed Tables. Retrieved from U.S. Department of Health and Human Services, Substance Abuse and Mental Health Services Administration: <https://www.samhsa.gov/data/sites/default/files/cbhsqreports/NSDUHDetailedTabs2017/NSDUHDetailedTabs2017.htm#lotsect1pe>
- U.S. Department of Health and Human Services, Substance Abuse and Mental Health Services Administration. (2018). Interagency Coordinating Committee on the Prevention of Underage Drinking. Retrieved from Report to Congress on the Prevention and Reduction of Underage Drinking: <https://www.stopalcoholabuse.gov/resources/reporttocongress/rtc2018.aspx>
- U.S. Department of Health and Human Services, Substance Abuse and Mental Health Services Administration. (2020, September 11). 2019 NSDUH Detailed Tables. Retrieved from 2019 NSDUH Detailed Tables: <https://www.samhsa.gov/data/report/2019-nsduh-detailed-tables>.
- U.S. Department of Health and Human Services, Substance Abuse and Mental Health Services Administration. (2021, March). Facts on College Drinking: March 2021. Retrieved from Substance Abuse and Mental Health Services Administration: https://store.samhsa.gov/sites/default/files/SAMHSA_Digital_Download/PEP21-03-10-006.pdf
- U.S. Department of Justice, Office of Justice Programs (2020, September). Promising Strategies to Reduce Substance Abuse. Retrieved April 16, 2023, from <https://www.ojp.gov/pdffiles1/ojp/183152.pdf>
- Varghese, M., Sheffer, C., Stitzer, M., Landes, R., Brackman, S., & Munn, T. (2014). Socioeconomic disparities in telephone-based treatment of tobacco dependence. *American Journal of Public Health*, e76-e84.
- Wang, T. W., Tynan, M. A., Hallett, C., Walpert, L., Hopkins, M., Konter, D., & King, B. A. (2018, June 21). Smoke-free and tobacco-free policies in colleges and Universities — United States and territories, 2017. Centers for Disease Control and Prevention. Retrieved April 12, 2023, from <https://www.cdc.gov/mmwr/volumes/67/wr/mm6724a4.htm>
- Webb Hooper, M., Antoni, M., Okuyemi, K., Dietz, N., & Rescinow, K. (2017). Randomized controlled trial of group-based culturally specific cognitive behavioral therapy. *Nitocine Tobacco Research*, 333-341.
- Webb Hooper, M., Carpenter, K., & Salmon, E. (2019). Web-based tobacco cessation interventions and digital inequality across US racial/ ethnic groups. *Ethnicity & Disease*, 495-504.
- Webb, M. (2009). Culturally specific interventions for African American smokers: An efficacy experiment. *Journal of the National Medical Association*, 101(9), 927-935.

- Wechsler, H., Lee, J., Kuo, M., & Lee, H. (2000). College binge drinking in the 1990s: a continuing problem, Results of the Harvard School of Public Health 1999 College Alcohol Study. *Journal of American College Health*, 199-210.
- Welsh, J. W., Shentu, Y., & Sarvey, D. B. (2019). Substance Use Among College Students. *Focus: Journal of Life Long Learning in Psychiatry*, 17(2), 117-127.
- White, A., Hingson, R., Pan, I.-J., & Hsiao-ye yi. (2011). Hospitalizations for Alcohol and Drug Overdoses in Young Adults Ages 18–24 in the United States, 1999–2008: Results From the Nationwide Inpatient Sample. *Journal of Studies on Alcohol and Drugs*, 72(5), 774-786.
- Wilson, O. W., Bullen, C., Duffey, M., & Bopp, M. (2022). The association between vaping and health behaviors among undergraduate college students in the United States. *Journal of American College Health*, 1360-1364.
- Witkiewitz, K., Desai, S., Bowen, S., Leigh, B., Kirouac, M., & Larimer, M. (2014). Development and evaluation of a mobile intervention for heavy drinking and smoking among college students. *Psychology of Addictive Behaviors*, 28(3), 639-650.
- Wolaver, A. (2002). Effects of heavy drinking in college on study effort, grade point average, and major choice. *Contemporary Economic Policy*, 20, 415-428.
- Zhu, S., Gardiner, P., Cummins, S., Anderson, C., Wong, S., Cowling, D., & Gamst, A. (2011). Quitline utilization rates of African-American and white smokers: The California experience. *American Journal of Health Promotion*, 25(5 Suppl), s51-s58.



©

UA LITTLE ROCK SURVEY RESEARCH CENTER

2801 South University Avenue | Ross Hall 404 | Little Rock, Arkansas 72204-1099 (O)

501.569.8559 | <https://ualr.edu/publicaffairs/survey-research-center/>